

PROVISIONAL APPLICATION
FILING RECEIPT



UNITED STATES PATENT AND TRADEMARK OFFICE
ASSISTANT SECRETARY AND COMMISSIONER
OF PATENTS AND TRADEMARKS
Washington, D.C. 20231

APPLICATION NUMBER	FILING DATE	FIL FEE REC'D	ATTORNEY DOCKET NO.	DRWGS
60/071,700	01/16/98	\$150.00	322029595	0

BARNES & THORNBURG
11 SOUTH MERIDIAN STREET
INDIANAPOLIS IN 46204

RECEIVED

MAR 26 1998

Receipt is acknowledged of this Provisional Application. This Provisional Application is not to be used for patentability. Be sure to provide the PROVISIONAL APPLICATION NUMBER, FILING DATE, NAME OF APPLICANT, and TITLE OF INVENTION when inquiring about this application. Fees transmitted by check or draft are subject to collection. Please verify the accuracy of the data presented on this receipt. If an error is noted on this Filing Receipt, please write to Exx Provisional Application within 10 days of receipt. Please provide a copy of the Provisional Application Filing Receipt with the changes noted thereon. This Provisional Application will automatically be abandoned twelve (12) months after its filing date and will not be subject to revival to restore it to pending status beyond a date which is after twelve (12) months from its filing date.

Applicant(s)

MARK CUSHMAN, WEST LAFAYETTE, IN; AGUSTIN
CASIMIRO-GARCIA, WEST LAFAYETTE, IN; WILLIAM RICE,
FREDRICK, MD.

FOREIGN FILING LICENSE GRANTED 03/20/90

TITLE

ALKENYLDIARYLMETHANE NON-NUCLEOSIDE HIV-1 REVERSE TRANSCRIPTASE
INHIBITORS

DOCKETED

DATE AUG 31, 1998

BY MDH

PROVISIONAL PATENT APPLICATION

of

Mark Cushman
(West Lafayette, Indiana)

Agustin Casimiro-Garcia
(West Lafayette, Indiana)

William G. Rice
(Fredrick, Maryland)

For

**ALKENYLDIARYLMETHANE NON-NUCLEOSIDE HIV-1 REVERSE
TRANSCRIPTASE INHIBITORS**

Attorney Docket 3220-29595

Attorneys:

John P. Breen
Steven R. Lammert
BARNES & THORNBURG
11 South Meridian Street
Indianapolis, Indiana 46204

**THE INVENTION DISCLOSED HEREIN WAS NOT MADE WITH GOVERNMENT
FUNDING TO CAUSE ASSIGNMENT TO THE GOVERNMENT.**

ALKENYLDIARYLMETHANE NON-NUCLEOSIDE HIV-1 REVERSE TRANSCRIPTASE INHIBITORS

Background

5 The non-nucleoside HIV-1 reverse transcriptase inhibitors (NNRTIs) are a structurally diverse set of compounds that inhibit reverse transcriptase by an allosteric mechanism involving binding to a site adjacent to the deoxyribonucleoside triphosphate binding site of the enzyme. Familiar examples of NNRTIs include hydroxyethoxymethylphenylthiothymine (HEPT), tetrahydroimidazobenzodiazepinone
10 (TIBO), dipyrindodiazepinone (nevirapine), pyridinone, bis(heteroaryl)piperazine (BHAP), *tert*butyldimethylsilylspiroaminooxathiole dioxide (TSAO), and α -anilinophenylacetamide (α -APA) derivatives. Nevirapine has recently been approved for clinical use as an anti-AIDS agent.

 The use of NNRTIs as anti-AIDs agents has been limited by the
15 development of viral resistance to the NNRTIs. Although the rapid emergence of resistant viral strains has hampered the clinical development of the NNRTIs for the treatment of AIDS, several strategies have emerged for overcoming resistance, including switching to another NNRTI to which the virus has remained sensitive, using higher doses of the NNRTI against the resistant strain, employing of combinations of agents which
20 elicit mutations that counteract one another, and combining NNRTIs with nucleoside reverse transcriptase inhibitors (RTIs). Accordingly, a need remains for additional NNRTIs having unique patterns of resistance mutations in order to facilitate the application of these strategies.

 The synthesis and biological evaluation of NNRTIs in the
25 alkenyldiarylmethane (ADAM) series has recently been reported. Several of the alkenyldiarylmethane compounds were disclosed as inhibiting the cytopathic effect of a wide variety of HIV-1 strains in CEM, MT-4, and monocyte-macrophage cultures. (Cushman et al. J. Med. Chem 1996, 39, 3217-3227) The most potent of these disclosed compounds was ADAM I (1), which displayed anti-HIV activity vs. a wide range of HIV-
30 1 isolates and was synergistic with AZT. However, the potency of ADAM I (1) against a variety of non-resistant HIV-1 strains was lower than that generally observed with many

of the known NNRTIs, ranging from 0.56 μM vs. HIV-1_{6S} in MT-4 cells to 151 μM vs. HIV-1_{NI19} in MT-4 cells.

The present invention is directed to a series of ADAM I related compounds that are inhibitors of reverse transcriptase activity and inhibit the cytopathic activity of HIV strains.

Detailed Description of the Invention


The design of additional alkenyldiarylmethanes has been aided by the availability of X-ray structures of HIV-1 reverse transcriptase complexed with nevirapine, α -APA, and TIBO. These structures reveal that nevirapine, α -APA, and TIBO assume a similar butterfly shape and bind to the enzyme in a similar manner with considerable overlap. Analysis of the X-ray crystallography structures allows the construction of a hypothetical model of the binding of ADAM I (1) to HIV-1 RT. The model was constructed by overlapping the structure of ADAM I (1) with that of nevirapine (2) in the binding pocket of HIV-1 RT (Sculpt™ 2.0, Interactive Simulations, San Diego, California). During this process, it was assumed that the hexenyl side chain of ADAM I (1) would point in the same direction as the cyclopropyl substituent of nevirapine. The nevirapine structure was then removed, the structure of the protein "frozen", and the energy of the complex minimized while allowing the ligand to move. The resulting hypothetical structure was consistent with the reported structures of NNRTI enzyme complexes, and was also supported by prior mutagenesis studies of the alkenyldiarylmethane binding site of HIV-1 reverse transcriptase, in which it was determined that the resistance mutations to ADAM I circumscribe a well-defined binding pocket.

According to the model generated by this analysis, the end of the ADAM I (1) side chain occupies a cavity formed by Glu 138 (top), Lys 103 (right), Tyr 181 (left), and Val 179 (bottom) of the HIV-1 RT. Several functional groups are present that would be capable of hydrogen bonding, including the phenolic hydroxyl group of Tyr 181, the backbone amide and side chain carboxylate of Glu 138, and the terminal amino group of Lys 103. It was anticipated that the incorporation of functional groups at the end of the alkenyl chain of the ligand which are capable of hydrogen bonding might allow favorable interactions with the adjacent residues of the RT. Therefore, alkenyldiarylmethane related


compounds were synthesized to incorporate functionalities at the end of the alkenyl side chain that would be capable of hydrogen bonding.

The present invention is directed to non-nucleoside compounds that inhibit reverse transcriptase activity, their pharmaceutical compositions and methods utilizing such compounds/compositions for treating patients suffering from a viral infection. More particularly the compounds of the present invention are useful for treating patients suffering from a disease of retroviral origin, such as AIDs.

The compounds of the present invention are alkenyldiarylmethane compounds of formula I:



wherein X is selected from the group consisting of



wherein R_1 and R_6 are H or halo;

R_2 and R_5 are independently OR_{11} ;

R_3 and R_4 are independently CO_2R_{12} ;

R_7 , R_8 , R_9 and R_{10} are independently selected from the group consisting of H, (C_1-C_5) alkyl, $(CH_2)_nOR_{13}$, $(CH_2)_nN_3$, $(CH_2)_nCOOR_{14}$, $(CH_2)_nOSO_2CH_3$ and $(CH_2)_nNH_3$;

R_{11} and R_{12} are independently (C_1-C_5) alkyl or H;

R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; and n is 0-4.

In one preferred embodiment of this invention there is provided compounds of the above formula I, wherein X is




R_1 and R_6 are independently Br or Cl, R_2 and R_5 are each OCH_3 , R_3 and R_4 are each CO_2CH_3 , R_7 is H and R_8 is $(\text{CH}_2)_m\text{OH}$, $(\text{CH}_2)_m\text{COOCH}_3$ or $(\text{CH}_2)_m\text{N}_3$, wherein m is 2 or 3. These compounds have been demonstrated to inhibit the cytopathic effect of HIV-1_{RF} in CEM-SS cells. (See Table 1).

In another preferred embodiment of this invention there is provided a reverse transcriptase inhibiting compounds of the above formula I wherein X is




R_1 and R_6 are halo;
 R_2 and R_5 are OCH_3 ;
 R_3 and R_4 are CO_2CH_3 ;
 R_7 is H;
 R_8 is $(\text{C}_2\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;
 R_{13} and R_{14} are independently selected from the group consisting of H and $(\text{C}_1\text{-C}_5)$ alkyl;
and m is 2-3.

The compounds of this invention are readily formulated into pharmaceutical compositions, also within the scope of this invention, for use in the presently described method for treatment of patients suffering from a disease of retroviral origin. In one preferred embodiment of this invention, the pharmaceutical composition comprises a reverse transcriptase inhibitory effective amount of an compound of formula I:



wherein X is selected from the group consisting of



- wherein R_1 and R_6 are H or halo;
 R_2 and R_5 are independently OR_{11} ;
 R_3 and R_4 are independently CO_2R_{12} ;
 10 R_7 and R_8 are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;
 R_9 and R_{10} are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_n\text{OR}_{13}$, $(\text{CH}_2)_n\text{N}_3$, $(\text{CH}_2)_n\text{COOR}_{14}$ and $(\text{CH}_2)_n\text{NH}_3$;
 R_{11} and R_{12} are independently $(\text{C}_1\text{-C}_5)$ alkyl or H;
 15 R_{13} and R_{14} are independently selected from the group consisting of H and $(\text{C}_1\text{-C}_5)$ alkyl;
 m is 1-3 and n is 0-4 and a pharmaceutically acceptable carrier.

Another pharmaceutical composition within the scope of this invention comprises a reverse transcriptase inhibiting compounds of the above formula I wherein X is



- R_1 and R_6 are halo;
 R_2 and R_5 are OCH_3 ;
 25 R_3 and R_4 are CO_2CH_3 ;
 R_8 is $(\text{C}_2\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;
 R_{13} and R_{14} are independently selected from the group consisting of H and $(\text{C}_1\text{-C}_5)$ alkyl;
 and m is 2-3 and a pharmaceutically acceptable carrier.

- The present invention further provides pharmaceutical formulations
 30 comprising an effective amount of an alkenyldiarylmethane compound for treating a patient suffering from a disease of retroviral origin. As used herein, an effective amount of the alkenyldiarylmethane compound is defined as the amount of the compound which,

upon administration to a patient, alleviates or eliminates symptoms of the disease, or reduces or eliminates detectable levels of the virus in the treated patient.

The effective amount to be administered to a patient is typically based on body surface area, patient weight, and patient condition. Body surface area may be approximately determined from patient height and weight (see e.g., Scientific Tables, Geigy Pharmaceuticals, Ardley, New York, pages 537-538 (1970)). Effective doses will vary, as recognized by those skilled in the art, dependant on route of administration, excipient usage and the possibility of co-usage with other therapeutic treatments including other anti-tumor agents, and radiation therapy.

The pharmaceutical formulation may be administered via the parenteral route, including subcutaneously, intraperitoneally, intramuscularly and intravenously. Examples of parenteral dosage forms include aqueous solutions of the active agent, in a isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carrier. In one preferred aspect of the present embodiment, the alkenyldiarylmethane compound is dissolved in a saline solution containing 5% of dimethyl sulfoxide and 10% Cremphor EL (Sigma Chemical Company). Additional solubilizing agents well-known to those familiar with the art, can be utilized as pharmaceutical excipients for delivery of the selenophene compounds.

The present compound can also be formulated into dosage forms for other routes of administration utilizing well-known methods. The pharmaceutical compositions can be formulated, for example, in dosage forms for oral administration in a capsule, a gel seal or a tablet. Capsules may comprise any well-known pharmaceutically acceptable material such as gelatin or cellulose derivatives. Tablets may be formulated in accordance with conventional procedure by compressing mixtures of the active compound and solid carriers, and lubricants well-known to those familiar with the art. Examples of solid carriers include starch, sugar, bentonite. The compounds of the present invention can also be administered in a form of a hard shell tablet or capsule containing, for example, lactose or mannitol as a binder and conventional fillers and tableting agents.

The anti-viral activity of the described compounds was measured utilizing two different assays. The first assay measures the effectiveness of the alkenyldiarylmethane compound to inhibit reverse transcriptase activity, and the second assay measured the compound's ability to inhibit the cytopathic activity of HIV-1 to cells

cultured *in vitro*. The mechanism of action for the disclosed compounds' antiviral activities is believed to be due at least in part to the compounds' ability to inhibit reverse transcriptase (RT) activity. Several of the disclosed ADAM compounds have an inhibitory effect on RT and yet do not exhibit a detectable inhibitory effect on viral cytopathicity in the HIV cell assay. This may be the result of the failure of the compound to penetrate the cell membrane. For example, compound 21 has activity as an RT inhibitor and yet no detectable inhibitory cytopathic effect was detected. The inactivity of 21 might possibly be due to the fact that the amino group is protonated at the pH of the assay medium and therefore should be less able to penetrate cellular membranes. Such compounds may have utility as antiviral agents, if they can be formulated as prodrugs, using techniques known to those skilled in the art, or through the use of other techniques that are known to increase cellular uptake of compounds.

The following examples are provided to illustrate various embodiments of Applicants' invention, and are not intended to in any way limit the scope of the invention as set forth in this specification and appended claims.

Example 1


Preparation of Congeners of compound 1

Congeners of 1 were prepared in which the effect of alkenyl chain length on activity could be investigated. Various analogs were also prepared in which the bromines present in 1 were replaced by chlorines, iodines, and hydrogens.

Considering first the compounds having the same substitution in the aromatic rings as in 1, compounds 3-7 were prepared from the intermediate benzophenone 10. The methylene and ethylene compounds 3 and 4 were prepared by the Wittig reactions of 10 starting from methyltriphenylphosphonium bromide and ethyltriphenylphosphonium bromide, respectively, using sodium bis(trimethylsilyl)amide as the base. The methoxyethylene congener 5 was prepared in a similar reaction employing methoxymethyl(triphenyl)phosphonium bromide. Similarly, reaction of ketone 10 with the Wittig reagent derived from 3-[*tert*-(butyldiphenylsilyloxy)propyl]triphenylphosphonium bromide afforded intermediate 6. Removal of the protecting group from 6 was accomplished with tetra-*n*-butylammonium fluoride in THF, yielding the alcohol 7. Reaction of the alcohol 7 with mesyl chloride in the presence of triethylamine


5

10




15


1



2



20




10 X = Br

11 X = Cl

- 3 R = H
 4 R = CH₃
 5 R = OCH₃
 6 R = CH₂CH₂OTBDPS
 7 R = CH₂CH₂OH
 8 R = CH₂CH₂OSO₂CH₃
 9 R = CH₂CH₂N₃

25



12

30

gave the mesylate 8, which was converted to the azide 9 on treatment with sodium azide in DMF.



In order to convert 1 into a closely related compound bearing a reactive functionality which might serve to alkylate the enzyme, the conversion of the alkene moiety to an epoxide was considered. Reaction of 1 with *m*-chloroperoxybenzoic acid in methylene chloride afforded the desired epoxide 12.

A series of dichloro alkenyldiarylmethanes 13-23 bearing alkenyl substituents capable of hydrogen bonding was prepared using the substituted dichlorobenzophenone 11 as the starting material. Using the Horner-Emmons reaction, compound 24 was reacted with NaH at low temperature for 1 h to produce the corresponding anion, which was then reacted with ketone 11 to afford, after purification, ester 13 in good yield. The required reagent 24 was prepared from diethylphosphonoacetic acid (25) and 2-(trimethylsilyl)ethanol (26). Deprotection of 13 was carried out with tetra-*n*-butylammonium fluoride in THF, affording the carboxylic acid 14 as a colorless crystalline solid. The methyl ester 15 and *tert*-butyl ester 16 were also prepared by the Horner-Emmons reaction starting from methyl diethylphosphonoacetate (27) and *tert*-butyl diethylphosphonoacetate (28), respectively.

The Wittig reaction of the ketone 11 with the ylide derived from 3-[*tert*-(butyldiphenylsilyloxy)propyl]triphenylphosphonium bromide afforded the alkene 17. Removal of the *tert*-butyldiphenylsilyl protecting group was accomplished with tetra-*n*-butylammonium fluoride in THF to provide the alcohol 18. Oxidation of 18 under Jones conditions (CrO₃, sulfuric acid, acetone), followed by extraction with 3 M aqueous sodium hydroxide, yielded the tricarboxylic acid 23, resulting from oxidation of the primary alcohol in 18 and hydrolysis of the two ester groups during extraction. On the other hand, reaction of the alcohol 18 with methanesulfonyl chloride in dichloromethane, using triethylamine as the base, afforded the mesylate 19. Reaction of the mesylate 19 with sodium azide in DMF gave the azide 20.

Various methods were considered for the reduction of the azide 20 to the amine 21. The usual methods for conversion of azides to amines involving lithium aluminum hydride or catalytic hydrogenations are obviously not suitable for accomplishing the conversion of 20 to 21 because of the other functionality in addition to the azide present in 20. Following a protocol described by Brown and Salunkhe,

5

24 R = CH₂CH₂TMS


25 R = H

27 R = CH₃28 R = C(CH₃)₃


10

13 R = COOCH₂CH₂TMS

14 R = COOH


15 R = COOCH₃16 R = COOC(CH₃)₃17 R = CH₂CH₂OTBDPS18 R = CH₂CH₂OH19 R = CH₂CH₂OSO₂CH₃20 R = CH₂CH₂N₃21 R = CH₂CH₂NH₂22 R = CH₂CH₂CH₂COOCH₃

26



29



15



23

20

25

30 R = CH₂CH₂OH31 R = CH₂CH₂Br

32

30

(Tetrahedron Lett. 1995, 36, 7987-7990) compound **20** was reacted with dichloroborane dimethyl sulfide complex to produce, after appropriate work up, amine **21** in low yield (15%). Dichloroborane dimethyl sulfide complex has been reported to react with olefins, but this process has been described to be slow and incomplete in the absence of trichloroborane. However, it is likely that the reactivity of the borane complex with olefins is at least in part responsible for the poor yield observed in the conversion of **20** to amine **21**. Alternative methods were then considered in order to reduce **20** to **21** in acceptable yield. Zwierzak et al. reported the preparation of amines from azides using a variant of the Staudinger reaction. (Synthesis, 1985, 202-204) They reacted organic azides with trialkyl phosphates, instead of the commonly used triphenylphosphine, to produce iminophosphorane intermediates, which after treatment with hydrogen chloride afforded the corresponding amine hydrochlorides. The higher reactivity of trialkyl phosphates towards azides was the main assumption for the modification. This methodology was considered to be the most adequate for the conversion of azide **20** to amine **21**, since the other reactive functionality present in **21** would not be affected by this method. In fact, the reaction of azide **20** with triethylphosphite for 24 h, followed by hydrolysis of the intermediate iminophosphorane with dry hydrogen chloride for 48 h, afforded the amine **21** as its hydrochloride salt in 86% yield.

In order to synthesize an alkenyldiarylmethane having a terminal methoxycarbonyl group at the end of an extended alkenyl side chain, the benzophenone **11** was subjected to a Wittig reaction with the ylide derived from the reaction of the phosphonium bromide **29** with sodium bis(nimethylsilyl)amide in THF. This resulted in the formation of the desired compound **22**, referred to herein as ADAM II

The effect of replacement of the methoxycarbonyl and methyl ether groups by cyclic acetonides on biological activity could be readily investigated by evaluating the known alcohol **30**, prepared previously from the ketone **32**. The primary alcohol was converted to the corresponding bromide **31** with carbon tetrabromide and triphenylphosphine in acetonitrile.

In order to determine the importance of the two halogen atoms for biological activity, congeners of ADAM **1** were prepared in which the halogens were replaced by iodines and by hydrogens. The syntheses of these compounds is outlined in Scheme 1. Treatment of the known diphenylmethane **33** with dimethylsulfate in refluxing

5

Scheme 1^a


10

15

20

25

30



^aReagents: (a) Me_2SO , K_2CO_3 , Me_2CO , reflux (6 h); (b) CrO_3 , Ac_2O , 0°C (1 h) and RT (12 h); (c) (1) $n\text{-C}_6\text{H}_{13}\text{PPh}_3\text{Br}$, $\text{NaN}(\text{SiMe}_3)_2$, THF, 0°C (30 min), (2) add 35, 23°C (12 h); (d) HNO_3 , Ac_2O , RT (12 h); (e) H_2 , PtO_2 , EtOAc ; (f) (1) NaONO , HCl , H_2O , 30 min (0°C), (2) I_2 , KI , H_2O , RT (30 min); (g) (1) $n\text{-C}_6\text{H}_{13}\text{PPh}_3\text{Br}$, $\text{NaN}(\text{SiMe}_3)_2$, THF, 0°C (30 min), (2) add 39, 23°C (12 h).

acetone with potassium carbonate as the base resulted in methylation of both carboxylic acids and both phenols to afford compound 34, which was oxidized to the benzophenone 35 with chromium trioxide in acetic anhydride. Reaction of the ketone present in 35 with the Wittig reagent derived from *n*-hexyltriphenylphosphonium bromide gave the desired analog 36, in which the two chlorines present in 1 have been replaced by hydrogens.

Nitration of 35 with nitric acid in acetic anhydride afforded the dinitro intermediate 37. The two nitro groups present in 37 were reduced to amines with Adam's catalyst in ethyl acetate to provide the diamino compound 38. Treatment of the diamine 38 with nitrous acid resulted in the conversion of both amines to diazonium groups, which were displaced by iodide in the presence of potassium iodide and iodine under aqueous conditions to provide intermediate 39. The reaction of 39 with the ylide derived from *n*-hexyltriphenylphosphonium bromide afforded the desired analog 40.

Example 2

Biological Results.

Nineteen new alkenyldiarylmethanes were tested for prevention of the cytopathic effect of HIV-1_{RF} in CEM-SS cells and for cytotoxicity in uninfected CEM-SS, and the results are listed in Table 1. In addition, they were tested as inhibitors of HIV-1 reverse transcriptase, and the resulting IC₅₀ values are also listed in Table 1. The most potent of the new alkenyldiarylmethanes, as well as the one with the highest therapeutic index, proved to be ADAM II (22), which displayed an EC₅₀ for prevention of the cytopathic effect of HIV-1_{RF} of 0.013 μ M. This represents an approximately 700-fold increase in potency over the lead compound 1, which was the most potent alkenyldiarylmethane of the previously disclosed ADAM series. Perhaps more importantly, the therapeutic index (the ratio of the CC₅₀ value to the EC₅₀ value) increased by a factor of 162. The other new compounds which were equipotent or more potent than 1 for the prevention of HIV-1 cytopathicity included the primary alcohol 7 (EC₅₀ = 6.0 μ M), the azide 9 (EC₅₀ = 1.1 μ M), the primary alcohol 18 (EC₅₀ = 8.6 μ M), and the azide 20 (EC₅₀ = 0.27 μ M). No inhibitory effect was detected for the remaining novel alkenyldiarylmethanes tested for inhibition of HIV-1 cytopathicity at concentrations up to those resulting in cytotoxicity in uninfected cells (see Table 1).

5

Table 1. Anti-HIV-1 Activities of ADAMs.

	compd	RT (IC ₅₀ μM) ^a	XTT Assay		TI ^d
			EC ₅₀ (μM) ^b	CC ₅₀ (μM) ^c	
10	1	0.38	9.2	138	15
	3	>100	NA ^e	316	-
	4	26	NA	10	-
	5	>100	NA	22	-
	7	31.6	6.0	>316	>52
	9	94	1.1	>316	>278
15	12	5.6	NA	>100	-
	14	>100	NA	48	-
	15	16	NA	25	-
	16	>100	NA	>100	-
	18	>100	8.6	16.8	2
	19	96	NA	6.9	-
	20	2.0	0.27	41.8	155
	21	63	NA	15	-
	22	0.3	0.013	31.6	2430
20	23	>100	NA	>316	-
	30	>100	NA	18.8	-
	31	>100	NA	176	-
	36	3.2	NA	14	-
	40	11	NA	>316	-

25

^aInhibitory activity vs. HIV-1 reverse transcriptase with rCdG as the template-primer. ^bThe EC₅₀ is the 50% inhibitory concentration for cytopathicity of HIV-1_{RF} in CEM-SS cells. ^cThe CC₅₀ is the 50% cytotoxic concentration for mock-infected CEM cells. ^dThe TI is the therapeutic index, which is the CC₅₀ divided by the EC₅₀. ^eNA means there was no observed inhibition of HIV-1 cytopathicity up to the cytotoxic concentration in uninfected cells.

30

The observed increase in antiviral potency of **22** relative to **1** did not correlate with inhibition of HIV-1 RT with poly(rC).oligo(dG) as the template primer, since **1** (IC_{50} vs. RT = 0.38 μ M) was actually more potent as an enzyme inhibitor than ADAM II (**22**) (IC_{50} vs. RT = 0.50 μ M). All of the new compounds displaying anti-HIV activity were also less potent than **1** as inhibitors of HIV-1 RT with poly(rC).oligo(dG) as the template primer, and some of them were significantly less active. Examples include compounds **7** (IC_{50} vs. RT = 31.6 μ M), **9** (IC_{50} vs. RT = 94 μ M), and **18** (IC_{50} vs. RT > 100 μ M). Conversely, analogs **12** and **36** inhibit HIV-1 RT with IC_{50} values of 5.6 μ M and 3.2 μ M, respectively, and are inactive as inhibitors of HIV-1 cytopathicity.


Some of the closely related structural analogs of **1** are revealing in their inactivity as inhibitors of the cytopathic effect of HIV-1. Examples include the epoxide **12**, the dechlorinated analog **36**, and the diiodo congener **40**. These inactive compounds emphasize the fact that there is a relatively high degree of structural specificity associated with the antiviral activity of the compounds in the series, so that even small changes in structure can result in complete loss of activity. The inactivity of **36** and **40** emphasize a significant role played by the two bromine atoms present in **1**. In this regard, the dichloro analog **41** was reported previously to retain activity, although it was approximately half as potent as **1**. However, the effect of bromine vs. chlorine substitution is not always consistent. For example, if one considers the bromo alcohol **7** (EC_{50} 6.0 μ M) vs. the chloro alcohol **18** (EC_{50} of 8.6 μ M), they are approximately equipotent as inhibitors of viral cytopathicity. On the other hand, the chloro azide **20** (EC_{50} 0.27 μ M) is four times more potent than the bromo azide **9** (EC_{50} 1.1 μ M).

The effect of replacement of the methoxycarbonyl and methyl ether groups by cyclic acetonides can be seen by comparison of the activities of the primary alcohols **18** and **30**. While **18** inhibited the cytopathic effect of HIV-1_{RF} with an EC_{50} of 8.6, the cyclic acetonide **30** was inactive. The corresponding primary bromide **31** was also inactive.

There appears to be an importance effect of chain length of the alkenyl appendage. Both of the active azides **9** and **20** have a chain length which is identical with the alkenyl chain present in ADAM **1**, and in the most active compound, ADAM II (**22**), the chain length is only slightly longer.

5

10



41

15

20

25

30

The effect of incorporation of hydrogen bonding groups at the end of the alkenyl chain seems to be variable, depending on the specific group incorporated and the length of the chain. For example, the primary alcohol **18** was effective in preventing the cytopathic effect of HIV-1_{RF} with an EC₅₀ of 8.6 μ M, but the corresponding amine **21** was inactive at 100 μ M. The most effective compounds for inhibition of viral cytopathicity were ADAM II (**22**), having a methyl ester at the end of the chain, and the two azides **9** and **20**. Although these compounds do have hydrogen bond accepting groups at the end of the alkenyl chain, it is difficult to attribute their greater antiviral activity to a greater affinity for the enzyme, because they are not more potent than **1** as enzyme inhibitors, at least with poly(rC).oligo(dG) as the template-primer.

In order to determine whether or not ADAM II was indeed acting as an NNRTI compound **22** was tested in a number of assays representative of important events in the replication cycle of HIV-1. These included (in addition to RT inhibition with poly(rA).oligo(dT) and poly(rC).oligo(dG) as template primers), assays for inhibition of HIV-1 attachment/fusion to target cells and the activities of HIV-1 integrase and protease enzymes. The effect of the compound on nucleocapsid protein was also investigated. Compound **22** did not have any significant effect on integrase, protease, or nucleocapsid protein. (See Table 2). However, it inhibited RT with either poly(rA).oligo(dT) or poly(rC).oligo(dG) as template primers. The greater sensitivity to inhibition with poly(rC).oligo(dG) as the template/primer is characteristic of the HIV-1-specific non-nucleoside reverse transcriptase inhibitors. Interestingly, ADAM I **1**, which had an IC₅₀ of 0.38 μ M with poly(rC).oligo(dG) as the template primer did not inhibit the enzyme with poly(rA).oligo(dT) as the template-primer. Thus there exists a clear distinction between the abilities of ADAM I and ADAM II to inhibit RT activity with the poly(rA).oligo(dT) template primer system.

In addition to the molecular target-based mechanistic assays, ADAM II (**22**) was also evaluated in a time course assay to determine the site of action of the compound during the early phase of HIV-1 replication, as well as with latently HIV-1 infected U1 cells to probe for any antiviral actions during the post-infective late phase of viral replication. The profile of inhibition of ADAM II corresponded to that of an NNRTI, in which antiviral activity was diminished when the compound was added to cultures four hours after initiation of infection. This is analogous to the effects of

nevirapine, but unlike the action of dextran sulfate, which blocks virus attachment at the cell surface and loses its effectiveness within the first half hour after initiation of infection. PCR-based analysis confirmed that ADAM II prevented formation of proviral DNA during the early phase of infection. ADAM II had no effect on virus production from U1
5 cells induced with TNF- α to produce virus from integrated proviral DNA. Thus, ADAM II acted biologically to inhibit replication by preventing reverse transcription.

The scope of the anti-HIV activity of ADAM II was investigated in a number of laboratory-adapted strains, lymphocyte-tropic clinical isolates, clade representatives, and monocyte-tropic strains of HIV-1. Likewise, ADAM II was tested
10 for inhibitory activity against an array of HIV-1 strains containing mutations in the reverse transcriptase enzyme. (See Tables 3, 4 and 5). The mutant viruses were obtained either by *in vitro* biological selection in the presence of various NNRTIs, or by site directed mutagenesis. In order to preserve continuity of data across the surfeit of virus strains, a greater amount of virus (higher MOI) was used in these competitive studies than was
15 used in the original screening assays. For this reason, the EC₅₀ was generally about 20-fold greater in these studies than in the original screen. Nevertheless, ADAM II retained potent anti-HIV-1 activity against a wide variety of laboratory and clinical HIV-1 strains in CEM-SS, PBMC, and Mono/Mac cells (Table 3). Results from studies with the panels of viruses having defined mutations in reverse transcriptase are listed in Tables 4 and 5.
20 Mutations conferring resistance to ADAM II clustered within the amino acid residues comprising the NNRTI binding pocket. Resistance mutations to the NNRTI α -APA are located close to the bound inhibitor, and it therefore seems likely that the mutations conferring resistance to ADAM II may also be located close to the bound ligand. The mutations conferring greater than ten-fold resistance to ADAM II were located at
25 positions 103, 108, 110, 139, 181, and 188. In contrast, the A98G and L100I mutants were more sensitive to ADAM II than the corresponding wild type virus (Table 5). The virus containing the L100I mutation and the multiple mutations that confer resistance to AZT also displayed increased sensitivity to ADAM II. This suggests the possible clinical utility of ADAM II against AZT resistant strains of HIV-1 and against strains of HIV-1
30 that express L100I-based resistance to other NNRTIs.

5

10

Table 2. Mechanistic Evaluations of ADAM II (22)

Parameter	IC ₅₀ (μM)
Attachment/Fusion	NI ₁₀₀ ^a
RT rAdT	1.9
rCdG	0.3
Integrase	NI ₁₀₀ ^b
Protease	>100 ^c
Nucleocapsin p7 Protein	NI ₂₅

15

^aA 20% inhibition was observed at 100 μM. ^bNI indicates that no inhibition of activity was observed at the indicated high test concentration. ^cA 30% inhibition of HIV-1 protease was observed at a concentration of 100 μM.

20

25

30

Table 3. ADAM II Activity Against HIV-1 Laboratory-Adapted Strains, Lymphocyte-Tropic Clinical Isolates, Clades and Monocyte-tropic Strains of HIV-1

Virus Isolate	Virus Type	Cell Type	EC ₅₀ (μM)
HIV-1 RF	WT-Laboratory ^a	CEM-SS	0.91
HIV-1 IIB	WT-Laboratory ^a	CEM-SS	0.16
HIV-1 TEKI	WT-Clinical ^b	PBMC	2.20
HIV-1 ROJO	WT-Clinical ^b	PBMC	2.28
HIV-1	Clade A	PBMC	1.09
HIV-1	Clade B	PBMC	0.43
HIV-1	Clade C	PBMC	1.04
HIV-1	Clade D	PBMC	1.02
HIV-1	Clade E	PBMC	0.40
HIV-1	Clade F	PBMC	1.70
HIV-1 Ba-L	Monocyte-tropic	Mono/Mac	0.36
HIV-1 ADA	Monocyte-tropic	Mono/Mac	0.28

^aWild type laboratory-adapted strain. ^bWild type clinical isolate.

Table 4. ADAM II Activity Against a Biological Panel of Resistant Isolates in CEM-CC Cells

Virus Isolate	Mutation in RT	EC ₅₀ (μM)	Fold Resistance
HIV-1 IIIB	Wild Type	0.39	
OC ^a	L100I	0.94	2.4
TSAO/Cost	K110E	11.9	30
129/Cost	K103N	10.8	28
Thiazol ^b	V108I	>50	>128
Calo ^c	T139I	15	38
DPS ^d	Y181C	>50	>128
3TC	M184I	1.08	2.8
Cost	Y188 H	>50	>128
HEPT ^e	P236L	8.08	21

^aOxathiin carboxanilide resistant. ^bThiazolobenzimidazole resistant.
^cCalanolide resistant. ^dDiphenyl sulfonate resistant.
^e(Hydroxyethoxy)methyl(phenylthio)thymine resistant.

Table 5. ADAM II Activity Against a Site Directed Panel of Resistant Isolates in CEM-SS Cells

Virus Isolate	Mutation in RT	EC ₅₀ (μM)	Fold Resistance
HIV-1 NL4-3	Wild type	1.78	
	Y188C	10.80	6.07
	K103N	>50	>28
	K101E	3.05	1.71
	L100I	<0.16	S ^a
	4XAZT/L100I ^b	0.27	S ^a
	A98G	0.84	S ^a
	V179D	50	28
	Y181C	>50	>28
	L74V	2.23	1.25
	4XAZT ^b	0.90	S ^a
	V108I	12.0	6.74
	V106A	12.7	7.13
	4XAZT/Y181C ^b	>50	>28

^aEnhanced Sensitivity. ^bAZT resistant.

Example 3

Experimental Section

General. Melting points were determined in capillary tubes on a Mel-Temp apparatus and are uncorrected. Spectra were obtained as follows: CI mass spectra on a Finnegan 4000 spectrometer, FAB mass spectra and EI mass spectra on a Kratos MS50 spectrometer, ¹H NMR spectra on Varian VXR-500S and Bruker ARX-300 spectrometers; IR spectra on a Beckman IR-33 spectrometer or on a Perkin Elmer 1600 series FTIR. Microanalyses were performed at the Purdue Microanalysis Laboratory, and all values were within $\pm 0.4\%$ of the calculated compositions.

3',3''-Dibromo-4',4''-dimethoxy-5',5''-bis(methoxycarbonyl)-1,1-diphenylethane (3). Methyl(triphenyl)phosphonium bromide (132 mg, 0.36 mmol) was suspended in anhydrous THF (2 mL, the mixture was stirred under Ar in an ice-bath, and sodium bis(trimethylsilyl)amide (1.0 M in THF, 0.4 mL, 0.4 mmol) was added by syringe. The mixture was stirred for 20 min, and a solution of 3,3'-dibromo-4,4'-dimethoxy-5,5'-bis(methoxycarbonyl)diphenyl ketone (10) (155 mg, 0.3 mmol) in anhydrous THF (3 mL) was slowly injected. The ice bath was removed, and the reaction mixture was stirred for 1 h at ambient temperature and 1 h at 60 °C overnight at ambient temperature. The reaction was quenched with saturated ammonium chloride solution (3 mL). The yellow organic layer was separated and the aqueous layer was extracted with benzene (2 x 5 mL). The combined organic extracts were washed with brine (2 mL) and dried (Na₂ SO₄). The solvent was evaporated and the residue was purified by flash chromatography on silica gel (6 g), eluting with hexane-ethyl acetate (6:1) to give **3** (62 mg 42%) as a colorless solid: mp 144 - 145 °C; ¹H NMR (CDCl₃, 300 MHz) δ 7.70 (m, 2 H), 7.68 (m, 2 H), 5.53 (s, 2 H), 4.01 (s, 3 H), 4.00 (s, 3 H), 3.97 (s, 3 H), 3.96 (s, 3 H); IR (KBr) 2943, 1730, 1472, 1434, 1246, 1208, 1086, 992, 797, 726 cm⁻¹; CIMS *m/z* (rel intensity) 516 (27), 515 (96), 514 (21), 512 (M⁺, 5), 486 (10), 485 (45), 484 (21), 483 (100). Anal. (C₂₀H₁₈Br₂O₆·1H₂O) C, H.

3',3''-Dibromo-4',4''-dimethoxy-5',5''-bis(methoxycarbonyl)-1,1-diphenylpropene (4). Ethyl(triphenyl)phosphonium bromide (134 mg, 0.36 mmol) was suspended in anhydrous THF 2 mL, and the mixture was stirred under Ar in an ice-bath. Sodium bis(trimethylsilyl)amide (1.0 M in THF, 0.5 mL) was added by syringe. The mixture was stirred for 20 min, and a solution of 3,3'-dibromo-4,4'-dimethoxy-5,5'-

bis(methoxycarbonyl)diphenyl ketone (10) (155 mg, 0.3 mmol) in anhydrous THF (3 mL) was slowly injected. The ice-bath was removed, and the reaction mixture was stirred for 24 h at ambient temperature. The reaction was quenched with saturated ammonium chloride solution (3 mL). The yellow organic layer was separated and the aqueous layer was extracted with benzene (2 x 5 mL). The combined organic extracts were washed with brine (2 mL) and dried (Na₂SO₄). The solvent was evaporated and the residue purified by flash chromatography on silica gel (6 g), eluting with hexane-ethyl acetate (6:1), to afford 4 (55 mg, 35%) as a colorless solid: MP 76 - 77 °C; ¹H NMR (CDCl₃, 300 MHz) δ 7.54 (m, 2 H), 7.51 (m, 1 H), 7.49 (m, 1 H), 6.18 (q, J = 7.2 Hz, 1 H), 3.99 (s, 3 H), 3.93 (s, 6 H), 3.92 (s, 3 H), 1.77 (d, J = 7.2 Hz, 3 H); IR (KBr) 2950, 1732, 1475, 1286, 1263, 1207, 997, 725 cm⁻¹; CIMS *m/z* (rel intensity) 531 (62), 530 (29), 529 (94), 528 (23), 527 (M+1, 57), 500 (12), 499 (53), 498 (24), 497 (100), 495 (53). Anal. (C₂₁H₂₀OBr₂O₆) C, H.

3', 3''-Dibromo-4',4''-dimethoxy-5',5''-bis(methoxycarbonyl)-1-methoxy2,2-diphenylethane (5). Methoxymethyl(triphenyl)phosphonium bromide (257 mg, 0.75 mmol) was suspended in anhydrous THF (4 mL), the mixture was stirred under Ar in an ice bath, and sodium bis(trimethylsilyl)amide (1.0 M in THF, 0.75 mL, 0.75 mmol) was added by syringe. The mixture was stirred for 20 min, and a solution of 3,3'-dibromo-4,4'-dimethoxy-5,5'-bis(methoxycarbonyl)diphenyl ketone (10) (258 mg, 0.5 mmol) in anhydrous THF (4 mL) was slowly injected. The ice bath was removed, and the reaction mixture was stirred for 24 h at ambient temperature, 3 h at 60 °C, and overnight at ambient temperature. The reaction was quenched with saturated ammonium chloride solution (5 mL). The yellow organic layer was separated and the aqueous layer was extracted with benzene (2 x 10 mL). The combined organic extracts were washed with brine (4 mL) and dried (Na₂SO₄). The solvent was evaporated and the residue purified by flash chromatography on silica gel (15 g), eluting with hexane-ethyl acetate 6:1, to give compound 5 as a colorless oil (76 mg, 28%) having an R_f of 0.15 on silica gel when hexane-ethyl acetate (6:1) was used as the solvent system: ¹H NMR (CDCl₃, 300 MHz) δ 7.78 (d, J = 2.1 Hz, 1 H), 7.74 (d, J = 2.1 Hz, 1 H), 7.61 (d, J = 2.2 Hz, 1 H), 7.58 (d, J = 2.2 Hz, 1 H), 6.51 (s, 1 H), 4.01 (s, 6 H), 3.99 (s, 3 H), 3.97 (s, 3 H), 3.89 (s, 3 H); IR (neat) 2949, 1733, 1636, 1475, 1436, 1289, 1255, 1208, 1124, 1081, 1049, 998 cm⁻¹. HRFABMS calcd for C₂₁H₂₁Br₂O₇, *m/z* 542.9654. Found *m/z* 542.9661.

3',3''-Dibromo-4',4''-dimethoxy-5',5'-bis(methoxycarbonyl)-1,1-diphenyl-4-[(*tert*-butyldiphenylsilyl)oxy]-1-butene (6). 3-[(*tert*-

(Butyldiphenylsilyloxy)propyl]triphenylphosphonium bromide (595 mg, 0.93 mmol) was suspended in anhydrous THF (6 mL) and the mixture stirred under Ar on an ice bath.

5 Sodium bis(trimethylsilyl)amide (1.0 M in THF, 1 mL) was added by syringe. The reaction mixture turned into a bright orange solution and was stirred in an ice bath for 30 min. A solution of the ketone 10 (320 mg, 0.62 mmol) dissolved in anhydrous THF (5 mL) was added and the reaction mixture was stirred at ambient temperature for 24 h. The mixture was quenched with saturated ammonium chloride solution (10 mL), followed by
10 ethyl acetate (10 mL). The organic layer was separated and the aqueous layer was extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were washed with brine (15 mL) and dried (Na_2SO_4). The solvent was evaporated and the yellow oily residue purified by flash chromatography on silica gel (20 g), eluting with hexane-ethyl acetate 3:1, to give 6 (284 mg, 58%) as a yellow oil: ^1H NMR (CDCl_3 , 300 MHz) δ 7.67 (d, J = 7.7 Hz, 4 H), 7.58 (m, 3 H), 7.51 (d, J = 2.1 Hz, 1 H), 7.40 (m, 6 H), 6.10 (t, J = 7.4 Hz, 1H), 4.03 (s, 3 H), 3.98 (s, 3 H), 3.95 (s, 3 H), 3.93 (s, 3 H) 3.79 (t, J = 6.1 Hz, 2 H), 2.39 (m, 2 H), 1.09 (s, 9 H); IR (neat) 2951, 2860, 1734, 1473, 1265, 1205, 998, 704 cm^{-1} . Anal. ($\text{C}_{38}\text{H}_{40}\text{Br}_2\text{O}_7\text{Si}$) C, H.

3',3''-Dibromo-4-hydroxy-4',4''dimethoxy-5',5'-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (7). Compound 6 (266 mg, 0.33 mmol) was dissolved in dry THF (7 mL) and the solution stirred at 0 °C under Ar. A 1.0 M solution of tetrabutylammonium fluoride in THF (0.7 mL, 0.7 mmol) was added. The solution turned yellow and was stirred at 0 °C for 5.5 h. Brine (10 mL) was added and the phases were separated. The aqueous phase was extracted with ethyl acetate (3 x 20 mL). The extract was dried
25 (MgSO_4) and the solvent was evaporated. The residue was purified by flash column chromatography on silica gel, eluting with hexane-ethyl acetate 3:1, 1:1 to give 7 (129 mg, 70%) as a yellowish oil: ^1H NMR (CDCl_3 , 300 MHz) δ 7.58 (m, 3 H), 7.52 (m, 1 H), 6.14 (t, J = 7.1 Hz, 1 H), 4.00 (s, 3 H), 3.96 (s, 3 H), 3.94 (s, 3 H), 3.93 (s, 3 H), 3.77 (t, J = 6.1 Hz, 2 H), 2.41 (q, J = 6.3 Hz, 2 H), 1.59 (brs, 1 H); IR (neat) 3430 (broad band),
30 2951, 1732, 1474, 1265, 1208, 998 cm^{-1} . Anal. ($\text{C}_{22}\text{H}_{22}\text{Br}_2\text{O}_7$) C, H.

3',3''-Dibromo-4-methanesulfonyloxy-4',4''dimethoxy-5',5'-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (8). Compound (7) (310 mg, 0.55 mmol)

and anhydrous triethylamine (0.23 mL, 1.7 mmol) were dissolved in dry dichloromethane (7 mL) and the mixture was stirred under Ar at 0 °C. Mesyl chloride (0.2 mL, 2.56 mmol) was added and the mixture was stirred at 0 °C for 3 h. The reaction mixture was diluted with dichloromethane (10 mL) and washed with 0.5 N HCl (2 x 20 mL, followed by sat NaHCO₃ (1 x 20 mL) and brine (1 x 20 mL). The organic extract was dried (MgSO₄). The solvent was evaporated and the residue purified on silica gel (16 g), eluting with hexane-ethyl acetate 3:1, to afford the mesylate **8** (270 mg, 77%) as a colorless oil: ¹H NMR (CDCl₃, 300 MHz) δ 7.55 (m, 2 H), 7.51 (m, 2 H), 6.14 (t, J = 7.4 Hz, 1 H), 4.30 (t, J = 6.2 Hz, 2 H), 3.99 (s, 3 H), 3.931 (s, 3 H), 3.927 (s, 3 H), 3.92 (s, 3 H), 3.04 (s, 3 H), 2.56 (q, J = 7.0 Hz, 2 H); IR (neat) 2952, 1732, 1474, 1358, 1265, 1175, 1089, 995 cm⁻¹. Anal. (C₂₃H₂₄Br₂SO₉) C, H.

4-Azido-3',3''-Dibromo-4',4''dimethoxy-5',5'-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (9). Compound **8** (231 mg, 0.363 mmol) was dissolved in dry *N,N*-dimethylformamide (5 mL). Sodium azide (120 mg, 1.82 mmol) was added and the mixture was stirred at 35-50 °C for 3 h. The reaction mixture was allowed to reach ambient temperature and then it was diluted with ethyl ether (42 mL). The ethereal solution was washed with water (2 x 35 mL), brine (1 x 35 mL), and dried (MgSO₄). The solvent was evaporated and the residue was purified on silica gel (16 g), eluting with hexane-ethyl acetate 3:1, to give **9** (132 mg, 62 %) as a colorless solid: mp 69 - 70 °C; ¹H NMR (CDCl₃, 300 MHz) δ 7.56 (m, 3 H), 7.50 (m, 1 H), 6.05 (m, 1 H), 3.99 (m, 1 H), 3.93 (m, 3 H), 3.40 (m, 2 H), 2.41 (m, 2 H); IR (KBr) 2944, 2105, 1731, 1473, 1436, 1246, 1207, 1085, 999, 806, 726 cm⁻¹. Anal. (C₂₂H₂₁Br₂N₃O₆) C, H.

3',3''-Dibromo-4',4''-dimethoxy-5',5''-bis(methoxycarbonyl)-1,1,diphenyl-1-heptene Epoxide (12). Compound **1** (67 mg, 0.116 mmol) was dissolved in dry methylene chloride (1 mL) and the solution cooled in a freezer for 5 min. 3-Chloroperoxybenzoic acid [minimum 57 % (70 mg)] was dissolved in CH₂Cl₂ (0.5 mL) and injected into the solution. The mixture was allowed to reach ambient temperature and was stirred for 21 h. Solvent was evaporated and the residue was purified by flash column chromatography on silica gel (7.5 g), eluting with hexane-ethyl acetate 20:1 followed by 10:1, to give the epoxide **12** (43 mg, 62 %) as an oil: ¹H NMR (CDCl₃, 300 MHz) δ 7.74 (m, 2 H), 7.67 (m, 1 H), 7.65 (m, 1H), 3.97 (s, 3 H), 3.94 (s, 3 H), 3.93 (s,

3 H), 3.92 (s, 3 H), 3.35 (m, 1 H), 1.45 (m, 4 H) 1.27 (m, 5 H), 0.88 (m, 3 H). IR (neat) 2953, 1732, 1472, 1435, 1260, 1207, 1087, 999, 720 cm^{-1} . Anal. ($\text{C}_{25}\text{H}_{28}\text{Br}_2\text{O}_7$) C, H.

2-(Trimethylsilyl)ethyl 3',3''-Dichloro-4',4''-dimethoxy-5,5''-bis(methoxycarbonyl)-3,3-diphenylpropenoate (13). A suspension of sodium hydride (0.018 g, 0.750 mmol) in dry THF (5 mL) was stirred in an ice bath under an argon atmosphere. Phosphonoacetate 24 (0.21 mL, 0.70 mmol) was added and the mixture was stirred at 0 °C for 1 h. The initial suspension turned into a clear solution within minutes. A solution of ketone 11 (0.200g, 0.469 mmol) in dry THF (4 mL) was then added dropwise. The ice bath was removed and the reaction mixture was stirred at rt for 30 h. The solvent was removed and the residue was taken up in cold water (30 mL) and ethyl ether (30 mL). The layers were separated and the aqueous one was extracted with ethyl ether (2 x 30 mL). The combined organic fractions were washed with brine (2 x 30 mL), dried over magnesium sulfate and filtered. The solvent was removed *in vacuo* to give a residue. After flash chromatography (SiO_2 , 35 g; column dimensions: 3 cm x 6.5 inch), eluting with hexanes:ethyl acetate 5:1, compound 13 (0.216 g, 81.2%) was obtained as a thick oil: ^1H NMR (300 MHz, CDCl_3) δ 7.60 (d, J = 2.4 Hz, 1H), 7.53 (d, J = 2.2 Hz, 1H), 7.39 (d, J = 3.1 Hz, 1 H), 7.38 (d, J = 2.5 Hz, 1 H), 6.31 (s, 1 H), 4.11 (m, J = 8.7 Hz, 2 H), 3.99 (s, 3 H), 3.95 (s, 3 H), 3.91 (s, 3 H), 3.90 (s, 3 H), 0.89 (m, J = 8.7 Hz, 2 H), 0.01 (s, 9 H); IR (film) 2952, 1737, 1477, 1251, 1161, 997, 859, 744 cm^{-1} ; FABMS (m/z): 568.8[MH] $^+$. Anal. ($\text{C}_{26}\text{H}_{30}\text{Cl}_2\text{O}_8\text{Si}$) C, H.

3',3''-Dichloro-4',4''-dimethoxy-5,5''-bis(methoxycarbonyl)-3,3-diphenylpropenoic Acid (14). A solution of ester 13 (0.198 g, 0.348 mmol) in dry THF (10 mL) was stirred under argon in an ice bath. A 1.0 M solution of TBAF (0.7 mL, 0.697 mmol) was added dropwise and the reaction mixture stirred at 0 °C for 1.5 h. The mixture turned into a light yellowish solution. Brine (30 mL) was added and the mixture stirred for 10 min. A 1.0 N solution of HCl (10 mL) was then added and the product was extracted with ethyl ether (3 x 30 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried over magnesium sulfate, filtered, and the solvent removed. Pure 14 (0.135 g, 83%) was obtained as a colorless crystalline solid after flash chromatography (SiO_2 , 20 g; column dimensions: 2.2 cm x 6.5 inch), using chloroform:methanol:formic acid (200:10:0.1 mL) as eluant: mp. 60-62 °C; IR (film) 3400-2900, 2955, 1731, 1477, 1262, 1210, 1164, 994, 743 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 7.59 (d, J = 2.4 Hz, 1

H), 7.50 (d, $J = 2.2$ Hz, 1 H), 7.37 (d, $J = 2.2$ Hz, 1 H), 7.35 (d, $J = 2.4$ Hz, 1 H), 6.29 (s, 1 H), 3.97 (s, 3 H), 3.93 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H); CIMS (m/z): 469 [MH, 25]⁺, 451 [MH-18, 100]⁺, 437 [MH-32, 23]⁺. Anal. (C₂₁H₁₈Cl₂O₈·0.4 H₂O) C, H.

- Methyl 3',3''-Dichloro-4',4''-dimethoxy-5,5''-bis(methoxycarbonyl)-3,3-diphenylpropenoate (15).** A suspension of sodium hydride (0.012 g, 0.493 mmol) in dry THF (5 mL) was stirred in an ice bath under an argon atmosphere. Methyl diethylphosphonoacetate (27) (0.09 mL, 0.469 mmol) was added and the mixture was stirred at 0 °C for 1 h. The initial suspension turned into a clear solution within minutes. A solution of ketone 11 (0.100 g, 0.235 mmol) in dry THF (5 mL) was added dropwise.
- 10 The ice bath was removed and the reaction mixture was stirred at rt for 30 h. Cold water (15 mL) was added and the mixture was stirred for 10 min. The product was extracted with ethyl ether (3 x 25 mL). The combined organic fractions were washed with brine (1 x 30 mL), dried over magnesium sulfate, and filtered. The solvent was removed *in vacuo* to give a residue. After purification by flash chromatography on silica gel (35g; column dimensions: 2 cm x 6 inch), eluting with hexanes:ethyl acetate 2:1, compound 15 (0.0948 g, 83.8%) was obtained as a thick colorless oil: ¹H NMR (300 MHz, CDCl₃) δ 7.59 (d, $J = 2.4$ Hz, 1 H), 7.53 (d, $J = 2.2$ Hz, 1 H), 7.37 (d, $J = 2.2$ Hz, 1 H), 7.36 (d, $J = 2.4$ Hz, 1 H), 6.32 (s, 1 H), 3.99 (s, 3 H), 3.94 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.63 (s, 3 H); IR (film) 2951, 1731, 1477, 1435, 1264, 1164, 996 cm⁻¹; CIMS (m/z): 483 [MH, 30]⁺ and 451 [(MH)⁺-CH₃OH, 100]. Anal. (C₂₂H₂₀Cl₂O₈) C, H.

- tert*-Butyl 3',3''-Dichloro-4',4''-dimethoxy-5,5''-bis(methoxycarbonyl)-3,3-diphenylpropenoate (16).** *tert*-Butyl diethylphosphonoacetate (28) (0.44 mL, 1.872 mmol) was dissolved in dry THF (15 mL) and the mixture was stirred under argon in an ice bath. A 1.0 M solution of sodium bis(trimethylsilyl)amide (2.0 mL, 2.01 mmol) was added and the solution was stirred at 0 °C for 1 h. At this time, a solution of the ketone 11³⁴ (0.400 g, 0.936 mmol) in dry THF (12 mL) was added dropwise and the resulting mixture was stirred at rt for 24 h. The solvent was evaporated and the residue partitioned between ethyl ether (40 mL) and water (60 mL). The layers were separated and the aqueous one was extracted with ethyl ether (2 x 40 mL). The combined organic extracts
- 25 were dried over magnesium sulfate, filtered, and the solvent removed *in vacuo* to give a yellowish oil. Purification by flash chromatography (SiO₂, 30 g), eluting with hexanes-ethyl acetate (3:1), afforded pure 10 (0.229 g, 46.6 %) as a white solid: mp 114-115.5

°C; ¹H-NMR (300 MHz, CDCl₃) δ 7.57 (d, *J* = 2.4 Hz, 1 H), 7.51 (d, *J* = 2.3 Hz, 1 H), 7.37 (d, *J* = 2.2 Hz, 1 H), 7.35 (d, *J* = 2.4 Hz, 1H), 6.22 (s, 3 H), 3.98 (s, 3 H), 3.94 (s, 3 H), 3.90 (s, 3 H), 3.88 (s, 3 H), 1.29 (s, 3 H); IR (film) 2952, 1731, 1479, 1368, 1258, 1095, 998, 848, 744 cm⁻¹; CIMS *m/z* (relative intensity) 451 (MH⁺-C₄H₈, 100), 469 (MH⁺-C₄H₈OH, 92), 470 (MH⁺-C₄H₈O, 64). Anal. (C₂₅H₂₆Cl₂O₈) C, H.

3', 3''-Dichloro-4', 4''-dimethoxy-5', 5'-bis(methoxycarbonyl)-1,1-diphenyl-4-[(*tert*-butyldiphenylsilyl)oxy]-1-butene (17). 3-[*tert*-(Butyldiphenylsilyloxy)propyl]triphenylphosphonium bromide (0.45 g, 0.702 mmol) was suspended in dry THF (5 mL) and stirred under argon. The suspension was cooled in an ice bath. A 1.0 M solution of NaHMDS (0.75 mL, 0.75 mmol) in THF was added dropwise. The reaction mixture turned into a bright orange solution and was stirred in the ice bath for 30 min. A solution of the ketone 11 (0.200 g, 0.468 mmol) in dry THF (2 mL) was added. The solution was stirred at room temperature for 24 h. A saturated solution of ammonium chloride (10 mL) was then added, followed by ethyl acetate (10 mL). The phases were separated and the aqueous one was extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried over magnesium sulfate, filtered, and the solvent evaporated to give a dark brown oil. Purification by flash chromatography (SiO₂), eluting with hexane/ethyl acetate (3:1), provided 17 (0.26 g, 76%) as a yellowish oil: ¹H-NMR (300 MHz, CDCl₃) δ 7.63 (dd, *J* = 1.4 Hz, 4 H), 7.49 (d, *J* = 2.3 Hz, 1H), 7.46 (d, *J* = 2.1 Hz, 1H), 7.31 (d, *J* = 2.0 Hz, 1 H), 7.28 (d, *J* = 2.3 Hz, 1H), 7.34 (m, 6 H), 6.10 (t, *J* = 7.4 Hz, 1H), 3.99 (s, 3 H), 3.94 (s, 3 H), 3.90 (s, 3 H), 3.88 (s, 3 H), 3.75 (t, *J* = 6.1 Hz, 2 H), 2.36 (t, *J* = 6.3 Hz, 1H), 2.34 (t, *J* = 6.4 Hz, 1 H), 1.05 (s, 9 H); IR (film) 3050, 2951, 1737, 1475, 1428, 1270, 1202, 1093, 999, 910, 702 cm⁻¹. Anal. (C₃₈H₄₀Cl₂O₇Si) C, H.

3', 3''-Dichloro-4-hydroxy-4', 4''dimethoxy-5', 5'-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (18). Silyl ether 17 (0.2 g, 0.282 mmol) was dissolved in dry THF (6 mL) and stirred at 0 °C under argon. A 1.0 M solution of tetrabutylammonium fluoride in THF (0.6 mL, 0.6 mmol) was added and the initially yellow solution turned into an orange solution which was stirred at 0 °C for 5.5 h. Brine (10 mL) was added and the phases were separated. The aqueous one was extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried over magnesium sulfate, filtered and the solvent evaporated to give an orange oil. Purification

by flash chromatography (silica gel) using hexane/ethyl acetate 1:1 as the eluant afforded 18 (94.5 mg, 71.4 %) as a colorless oil: ¹H NMR (300 MHz, CDCl₃) δ 7.52 (d, *J* = 2.4 Hz, 1 H), 7.50 (d, *J* = 2.1 Hz, 1H), 7.37 (d, *J* = 2.1 Hz, 1H), 7.31 (d, *J* = 1.8 Hz, 1H), 6.13 (t, *J* = 7.4 Hz, 1 H), 3.99 (s, 3 H), 3.93 (s, 3 H), 3.92 (s, 3 H), 3.91 (s, 3 H), 3.75 (t, *J* = 6.3 Hz, 2 H), 2.38 (q, *J* = 13.4 and *J* = 6.7 Hz, 1 H), 1.66 (bs, 1 H); IR (film) 3542, 2951, 2874, 1732, 1477, 1428, 1268, 1210, 1093, 998, 866, 743 cm⁻¹.

3', 3'' -Dichloro-4', 4'' -dimethoxy-5', 5'' -bis(carboxy)-4, 4'-diphenyl-3-butenic Acid (23). Chromium trioxide (0.64 g, 6.39 mmol) was dissolved in 1.5 M sulfuric acid (9.3 mL, 13.9 mmol), and the solution was stirred in an ice bath. A solution of alcohol 18 (0.50, 1.065 mmol) in acetone (15 mL) was then added and the ice bath removed. The reaction mixture was stirred at room temperature for 7 h. Ethyl ether (50 mL) was added and the phases separated. The organic phase was washed with water (3 x 40 mL) and then extracted with 3 M sodium hydroxide (3 x 20 mL). The combined aqueous extracts were acidified with conc hydrochloric acid and the cloudy solution was kept in the refrigerator overnight. Separation of the precipitated solid by filtration, followed by washing with water, afforded a yellowish solid (0.131 g), which was purified by recrystallization from ethyl ether/dichloromethane to yield 19 (63 mg, 12.2%) as a pale yellow solid: mp 206-207 °C; ¹H NMR (300 MHz, acetone-d₆) δ 7.65 (d, *J* = 2.3 Hz, 1 H), 7.62 (d, *J* = 2.1 Hz, 1 H), 7.57 (d, *J* = 2.1 Hz, 1 H), 7.52 (d, *J* = 2.3 Hz, 1 H), 6.42 (t, *J* = 7.4 Hz, 1 H), 3.97 (s, 6 H), 3.91 (s, 6 H), 3.20 (d, *J* = 7.4 Hz, 2 H); IR (film) 3400-2800, 2938, 1702, 1478, 1259, 996, 708 cm⁻¹; FABMS *m/z* (rel intensity): 454 (M⁺, 45), 437 (M⁺-17, 100). Anal. (C₂₀H₁₆Cl₂O₈·0.4 H₂O) C, H.

3', 3''-Dichloro-4-methanesulfonyloxy-4', 4''dimethoxy-5', 5''-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (19). A solution of alcohol 18 (0.700 g, 1.491 mmol) and triethylamine (0.62 mL, 4.473 mmol) in dry dichloromethane (20 mL) was stirred under argon at 0 °C. Mesyl chloride (0.35 mL, 4.473 mmol) was added and the mixture was stirred at 0 °C for 3 h. The reaction mixture was then diluted with dichloromethane (30 mL) and washed with 0.5 N HCl (2 x 40 mL), followed by sat NaHCO₃ (40 mL) and brine (40 mL). The organic extract was dried over magnesium sulfate, filtered, and the solvent removed *in vacuo* to produce a thick yellow oil (0.77 g). Purification of a fraction of this oil (0.557 g) by flash chromatography (25 g SiO₂) afforded pure 19 (0.521 g, 88.5 %) as a thick oil, which slowly crystallized. The

analytical sample was obtained by recrystallization from chloroform-methanol: mp. 83-85 °C; IR (film) 2952, 1731, 1479, 1359, 1269, 1174, 995 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3) δ 7.49 (d, $J = 2.4$ Hz, 1 H), 7.47 (d, $J = 2.2$ Hz, 1 H), 7.31 (d, $J = 2.1$ Hz, 1 H), 7.29 (d, $J = 2.3$ Hz, 1 H), 6.04 (t, $J = 7.4$ Hz, 1 H), 4.27 (t, $J = 6.3$ Hz, 2 H), 3.97 (s, 3 H), 3.91 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.00 (s, 3 H), 2.53 (q, $J = 6.4$ Hz and $J = 7.2$ Hz, 2 H); CIMS m/z (relative intensity) 547 (MH^+ , 46), 515 (100). Anal. ($\text{C}_{23}\text{H}_{24}\text{Cl}_2\text{O}_9\text{S}$) C, H.

4-Azido-3',3''-Dichloro-4',4''dimethoxy-5',5'-bis(methoxycarbonyl)-1,1-diphenyl-1-butene (20). Mesylate 19 (0.215 g, 0.393 mmol) was dissolved in dry DMF (5 mL). Sodium azide (0.13 g, 1.965 mmol) was added and the mixture was stirred at 35 to 50 °C for 3 h. The mixture was allowed to cool at room temperature, and then it was diluted with ether (45 mL). The ethereal solution was washed with water (2 x 40 mL), brine (1 x 40 mL), and dried over magnesium sulfate. After filtration and evaporation of the solvent *in vacuo*, a yellowish oil was obtained. Purification by flash chromatography (SiO_2 , 16 g), eluting with hexane:ethyl acetate 3:1, afforded pure 20 (0.148 g, 76.1 %) as an oil which slowly crystallized. The analytical sample was obtained by recrystallization from chloroform-pentane: mp 70-72 °C; ^1H NMR (300 MHz, CDCl_3) δ 7.50 (t, $J = 2.1$ Hz, 2 H), 7.34 (d, $J = 2.2$ Hz, 1 H), 7.30 (d, $J = 2.4$ Hz, 1 H), 6.05 (t, $J = 7.4$ Hz, 1 H), 3.99 (s, 3 H), 3.92 (s, 3 H), 3.91 (s, 3 H), 3.90 (s, 3 H), 3.39 (t, $J = 6.6$ Hz, 2 H), 2.39 (q, $J = 6.7$ Hz and $J = 7.2$ Hz, 2 H); IR (film) 2952, 2098, 1734, 1477, 1267, 997, 742 cm^{-1} ; CIMS m/z (relative intensity) 494 (MH^+ , 100), 496 (70). Anal. ($\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{N}_3\text{O}_6 \cdot 0.5 \text{H}_2\text{O}$) C, H.

4-Amino-3',3''-Dichloro-4',4'' dimethoxy-5',5'-bis(methoxycarbonyl)- 1,1-diphenyl-1-butene (21). A solution of azide 20 (0.228 g, 0.462 mmol) and $(\text{EtO})_3\text{P}$ (0.24 mL, 1.387 mmol) in benzene (6 mL) was stirred under argon at rt for 24 h. The reaction mixture was then saturated with dry HCl for 10 min and stirred for 48 h at rt. The solvent was removed and dry ethyl ether (15 mL) was added. The solution was placed in the freezer for 48 h. The precipitated amine hydrochloride was separated by filtration, washed with cold ethyl ether, and dried under high vacuum overnight to provide 21 (0.200 g, 86 % as a white solid. The analytical sample was recrystallized from ethanol-ethyl ether-pentane: mp 156-158 °C; ^1H NMR (300 MHz, CDCl_3) δ 7.49 (t, $J = 2.3$ Hz, 2 H), 7.47 (d, $J = 2.2$ Hz, 1 H), 7.32 (d, $J = 2.2$ Hz, 1 H), 7.29 (d, $J = 2.4$ Hz, 1 H), 6.06 (t, $J = 7.4$

Hz, 1 H), 3.96 (s, 3 H), 3.90 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3H), 2.81(t, $J=6.6$ Hz, 2H), 2.25 (q, $J=7.1$ Hz, 2H), 1.90 (bs, 2H, exchangeable with D_2O); IR (film) 2951, 1731, 1476, 1435, 1261, 1208, 997, 742 cm^{-1} ; CIMS m/z (rel intensity) 467 (M^+ , 90), 466 (M^+-1 , 100). Anal. ($C_{22}H_{23}Cl_2NO_6HCl$) C, H, N.

5 **Methyl 3',3''-Dichloro-4',4''-dimethoxy-5,5''-bis(methoxycarbonyl)-6,6-diphenylhexenoate (22).** (4-Methoxycarbonylbutyl)triphenylphosphonium bromide (29) (0.321 g, 0.704 mmol) was stirred in dry THF (15 mL) under argon at $-78^\circ C$. A 1.0 M solution of $NaN(SiMe_3)_2$ in THF (0.78 mL, 0.78 mmol) was then added and the yellow solution was stirred in a dry ice-acetone bath for 1 h. A $-78^\circ C$ solution of ketone 11
10 (0.200 g, 0.469 mmol) in dry THF (5 mL) was added and the reaction mixture was stirred at $-78^\circ C$ for 12 h, and then at rt for 12 h. Saturated NH_4Cl (25 mL) was added and the mixture stirred for 15 min. The layers were separated and the aqueous phase was extracted with ethyl acetate (3 x 30 mL). The combined organic extracts were washed with brine (1 x 40 mL), dried over magnesium sulfate, filtered, and the solvent removed *in vacuo* to give a thick orange residue. Purification was achieved by silica gel flash
15 chromatography, eluting with hexane:ethyl acetate 3:1, to provide 22 (0.110 g, 44.6 %) as a pale yellow oil: 1H NMR (300 MHz, $CDCl_3$) δ 7.51 (d, $J = 2.3$ Hz, 1 H), 7.48 (d, $J = 2.2$ Hz, 1 H), 7.33 (d, $J = 2.2$ Hz, 1 H), 7.30 (d, $J = 2.3$ Hz, 1 H), 6.05 (t, $J = 7.5$ Hz, 1 H), 4.01 (s, 3 H), 3.95 (s, 3 H), 3.94 (s, 3 H), 3.93 (s, 3 H), 3.65 (s, 3H), 2.32 (t, $J = 7.4$
20 Hz, 2 H), 2.15 (q, $J = 7.4$ Hz, 2 H), 1.80 (m, $J = 7.3$ Hz, 2 H); IR (film) 2951, 1736, 1477, 1261, 1208, 1093, 999, 743 cm^{-1} ; FABMS m/z : (rel intensity) 525 (MH^+ , 30), 509 (M^+-CH_3 , 28), 493 (M^+-OCH_3 , 100). Anal. ($C_{25}H_{26}Cl_2O_8$) C, H.

Following a similar procedure using 4-carboxybutyltriphenyl phosphonium bromide instead of 29 above, provided the corresponding 6,6-diphenyl hexanoic acid.

25 **(2-Trimethylsilyl)ethyl Diethylphosphonoacetate (24).** BOP-Cl (2.53 g, 9.952 mmol) was added to a solution of diethylphosphonoacetic acid (25) (1.6 mL, 9.952 mmol), (2-trimethylsilyl)ethanol (26) (1.57 mL, 10.94 mmol), and triethylamine (2.77 mL, 19.904 mmol) in dry dichloromethane (25 mL). The initial suspension turned into a clear solution within minutes, which was stirred at rt under Ar for 1.2 h. Water (60 mL, basified
30 with sodium bicarbonate) was added and the phases were separated. The organic phase was diluted with dichloromethane (2 x 30 mL) and washed with water (1 x 50 mL and brine (1 x 50 mL). The organic phase was dried over magnesium sulfate, filtered and the

solvent removed in vacuo to give a liquid residue (6.89 g). Purification by flash chromatography (SiO₂, 135 g) afforded **24** as a colorless liquid (2.47 g, 84%): bp 120-122 °C/ 0.05 mm Hg (Lit.³⁸ 140-145 °C/0.1 mm Hg); IR (film) 2981, 2954, 2904, 1737, 1268, 1027, 969, 838 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 4.18 (m, *J* = 8.6 Hz, 2 H), 4.14 (m, *J* = 7.2 Hz, 4 H), 2.94 (s, 1 H), 2.87 (s, 1 H), 1.31 (t, *J* = 7.1 Hz, 6 H), 0.98 (m, *J* = 8.7 Hz, 2 H), 0.01 (s, 9 H). Anal. (C₁₁H₂₅O₃PSi) C, H.

1-Bromo-4,4-bis(8',8''-dichloro-2',2'',2'',2''-tetramethyl-4',4''-dioxo-6',6''-0(1,3-benzodioxyl)]-3-butene (31). A solution of the alcohol **31** (0.106 g, 0.215 mmol) and carbon tetrabromide (0.09 g, 0.271 mmol) in dry acetonitrile (8.5 mL) was stirred under argon. A solution of triphenylphosphine (0.08 g, 0.305 mmol) in dry acetonitrile (1.5 mL) was added dropwise. The reaction mixture was stirred at reflux for 20 h. The solvent was removed *in vacuo* and the resulting oil was extracted with ethyl ether (5 x 5 mL). The solvent was evaporated from the combined organic extracts to give a thick solid. Purification by flash chromatography (SiO₂, 25 g), eluting with hexane-ethyl acetate 5:1, afforded the product **31** (66 mg, 56%) as a yellowish solid: mp 175-178 °C; ¹H NMR (CDCl₃, 300 MHz) δ 7.70 (d, *J* = 2.0 Hz, 1 H), 7.69 (d, *J* = 1.8 Hz, 1 H), 7.45 (d, *J* = 2.2 Hz, 1 H), 7.44 (d, *J* = 2.1 Hz, 1 H), 6.09 (t, *J* = 7.2 Hz, 1 H), 3.45 (t, *J* = 6.5 Hz, 2 H), 2.70 (q, *J* = 6.7 Hz, 2 H), 1.84 (s, 6 H), 1.80 (s, 6 H); IR (film) 2999, 1745, 1607, 1483, 1283, 1199, 1063, 874, 756 cm⁻¹; CIMS *m/z* (rel intensity) 555 (MH⁺, 55), 557 (MH⁺+2, 100), 559 (MH⁺+4, 40). Anal. (C₂₄H₂₁BrCl₂O₆) C, H.

4,4'-Dimethoxy-3,3'-bis(methoxycarbonyl)diphenylmethane (34). A suspension of 3,3'-dicarboxy-4,4'-dihydroxydiphenylmethane (**33**) (0.576 g, 2 mmol), dimethylsulfate (2 mL, 12 mmol), and potassium carbonate (2.0 g) in acetone (20 mL) was heated at reflux for 6 h. The solid mass was separated by filtration and washed with acetone. Evaporation of the acetone gave the product **34** (0.51g, 80%) as a brown oil: ¹H NMR (CDCl₃, 300 MHz) δ 7.61 (d, *J* = 1.8 Hz, 2 H), 7.25 (dd, *J* = 1.8, 8.6 Hz, 2 H), 6.90 (d, *J* = 8.5 Hz, 2H), 3.90 (s, 2 H), 3.87 (s, 6 H). Anal. (C₁₉H₂₀O₆) C, H.

4,4'-Dimethoxy-3,3'-bis(methoxycarbonyl)benzophenone (35). A solution of intermediate **34** (0.5 g, 1.8 mmol) in acetic anhydride (25 mL) was cooled to 0 °C in an ice bath containing NaCl. Solid chromium (VI) oxide (3 g, 30 mmol) was added slowly to the solution at 0 °C. After complete addition, the mixture was stirred at 0 °C for 1 h, and then room temperature for 12 h. The resulting viscous paste was broken up with ethyl

acetate and partitioned between 1 N HCl (200 mL) and ethyl acetate (200 mL). The organic layer was washed with 1 N HCl (100 mL) and brine (100 mL), dried over Na_2SO_4 , and the solvent removed under reduced pressure. Trituration of the solid in 1 N HCl and filtering gave the product 35 (0.12 g, 22%) as an off white solid: mp 142-144 °C; ¹H NMR (acetone-*d*₆) 8.14 (d, *J* = 2.2 Hz, 2 H), 7.96 (dd, *J* = 2.2, 8.7 Hz, 1 H), 7.30 (d, *J* = 8.7 Hz, 1 H), 3.99 (s, 3 H), 3.84 (s, 3 H). FTIR (KBr) 2953, 2849, 1735, 1710, 1650, 1603, 1502, 1438, 1406, 1274, 1152, 1081, 1010, 950 cm⁻¹. Anal. (C₁₉H₁₈O₇) C, H.

4',4''-Dimethoxy-3',3''-bis(methoxycarbonyl)- 1,1-diphenyl-1-heptene

(36). *n*-Hexyltriphenylphosphonium bromide (0.427 g, 1 mmol) was dried by azeotropic distillation from a benzene solution and then stirred in dry THF (10 mL) under nitrogen atmosphere. Sodium bis(trimethylsilyl)amide (1 M in THF, 1 mL, 1 mmol) was added and the ylide produced was stirred under nitrogen at 0 °C for 30 min. Intermediate 35 (0.276 g, 0.77 mmol) was added as a solution in THF (10 mL) under nitrogen. The mixture was stirred at rt overnight and partitioned between 1 N HCl (100 mL) and ethyl acetate (100 mL). The organic layer was dried over sodium sulfate, and evaporated to afford an oil which was chromatographed on SiO₂ (230-400 mesh, 50 g), eluting with hexanes/ethyl acetate (4:1) to give the product (36) (0.15 g, 46 %) as an oil: ¹H NMR (CDCl₃, 300 MHz) δ 7.63 (d, *J* = 2.3 Hz, 1 H), 7.56 (d, *J* = 2.2 Hz, 1 H), 7.20 (dt, *J* = 2.3 Hz, 1 H), 6.94 (d, *J* = 8.6 Hz, 1 H), 6.82 (d, *J* = 8.7 Hz, 1 H), 5.96 (t, *J* = 7.5 Hz, 1 H), 3.90 (s, 3 H), 3.84 (s, 3 H), 3.83 (s, 6 H), 2.05 (q, *J* = 7.3 Hz, 2 H), 1.39 (t, *J* = 7.2 Hz, 2 H), 1.22 (m, *J* = 1.9 Hz, 2 H), 0.83 (t, *J* = 6.7 Hz, 3 H); IR (neat) 2927, 2853, 1732, 1606, 1500, 1435, 1263 cm⁻¹. Anal. (C₂₅H₃₀O₆) C, H.

4,4'-Dimethoxy-3,3'-bis(methoxycarbonyl)-5,5'-dinitrobenzophenone (37).

A solution of compound 36 (1.07 g, 3 mmol) in acetic anhydride (30 mL) was cooled to 0 °C. Nitric acid 90 %, 20 mL was added dropwise and the solution was stirred overnight while warming to room temperature. The orange solution was poured onto ice and water and extracted with ethyl acetate (3 x 100 mL). The organic layer was washed with 5 % KOH solution (3 x 50 mL), dried over MgSO₄, and evaporated. The oil (1.1 g) was flash chromatographed on silica gel (250 g, 230-400 mesh), eluting with hexanes-ethyl acetate (10:2), to afford 37 (0.41 g, 32%) as a solid: mp 96-98 °C; ¹H NMR (CDCl₃, 300 MHz) 8.55 (s, 2 H), 8.41 (s, 2 H), 4.1 (s, 6 H), 3.99 (s, 6 H). Anal. (C₂₆H₁₆H₂O₇) C, H, N.

3, 3'-Diamino- 5,5'-Bis(methoxycarbonyl)-4,4'-dinitroxybenzophenone (38).

The dinitro compound 37 (0.33 g, 0.9 mmol) was hydrogenated at atmospheric pressure over platinum oxide (0.2 g, 0.08 mmol) in ethyl acetate (50 mL). After tlc (SiO₂, hexanes-acetone, 10:2) had shown that all the starting material was consumed, the catalyst was removed by filtration and the solvent was removed at reduced pressure to give an oil. This oil was flash chromatographed on SiO₂ (30 g, 230-400 mesh) using ethyl acetate. Evaporation of the solvent gave the product 38 (0.24 g, 80 %) as a glassy solid; ¹H NMR (CDCl₃, 300 MHz) δ 7.55 (d, J = 2.1 Hz, 2 H), 7.31 (d, J = 2.1 Hz, 2 H), 3.91 (s, 6 H), 3.89 (s, 6 H); IR (KBr) 3369, 2945, 2837, 1717, 1616 cm⁻¹. Anal. (C₁₉H₂₀N₂O₇)

3,3'-Diiodo-4,4',-dimethoxy-5,5'-bis(methoxycarbonyl)benzophenone (39). A

suspension of compound 38 (0.57 g, 1.4 mmol) in water (10 mL) was cooled to 0 °C. Concentrated HCl (0.6 mL) was added dropwise to give a yellow solution. After 10 min, sodium nitrite (0.22 g, 3.2 mmol) dissolved in water (2 mL) was added and the solution was stirred for 30 min at 0 °. The solution was then poured into a solution of iodine (1 g, 3.9 mmol) and potassium iodide (1.0 g, 5.9 mmol) in water (100 mL). The solution was stirred at room temperature for 30 min. Extraction with ethyl acetate (200 mL), washing with 10 % sodium hydrosulfate solution (100 mL), drying over MgSO₄, and evaporation of the solvent gave the crude diiodide (0.6 g). Recrystallization from hexanes/methylene chloride gave the pure product 39 (0.54 g, 63 %) as a solid: mp 160-161 °C. ¹H NMR (300 MHz, CDCl₃) δ 8.37 (d, J = 2.5 Hz, 2 H), 8.27 (d, J = 1.86 Hz, 2 H), 4.07 (s, 6 H), 3.95 (s, 6 H); IR (neat) 3078, 2954, 1734, 1670, 1608, 1540 cm⁻¹. Anal. (C₁₉H₁₆O₇I₂) C, H.

3',3'' -Diiodo-4',4''-Dimethoxy-5', 5''- bis(methoxycarbonyl)- 1, 1 diphenyl-

1-heptene (40). NaN(TMS)₂ (1 mL, 1 mmol) was added to an ice cold suspension of hexyltriphenylphosphonium bromide (0.427 g, 1 mmol) in THF (20 mL) and the solution was stirred for 30 min at 0 °C. A solution of the benzophenone 39 (0.2 g, 0.33 mmol) in THF (5 mL) was added to the preformed ylide and the solution was stirred at rt overnight. The mixture was partitioned between 1 N HCl (100 mL) and ethyl acetate (100 mL). The organic layer was evaporated and flash chromatographed on SiO₂ (230-400 mesh, 50.0 g), eluting with hexanes-ethyl acetate (5:1). The fractions which contained the product were pooled, evaporated, and rechromatographed on silica gel (230-400 mesh, 20 g), eluting with hexanes-ethyl acetate (5:1), to give the product 40

(0.022 g, 9 %) as an oil: ^1H NMR (CDCl_3 , 300 MHz) δ 7.69 (s, 2 H), 7.54 (d, $J = 2.0$ Hz, 1 H), 7.52 (d, $J = 2$ Hz, 1 H), 6.01 (t, $J = 7.6$ Hz, 1 H), 3.93 (s, 3 H), 3.89 (s, 3 H), 3.88 (s, 3 H), 3.85 (s, 3 H), 2.05 (q, $J = 7.3$ Hz, 2 H), 1.42 (m, 2 H), 1.25 (m, 2 H), 0.85 (t, $J = 6.7$ Hz, 3 H); IR (neat) 2953, 2929, 1742, 1738, 1731, 1713 cm^{-1} ; HRFABMS
 5 calcd for $\text{C}_{25}\text{H}_{28}\text{I}_2\text{O}_6$ m/z 677.9975 (M^+). Found: m/z 677.9954. Anal.
 ($\text{C}_{25}\text{H}_{28}\text{I}_2\text{O}_6 \cdot 0.5\text{EtOAc}$), C, H.

In Vitro Anti-HIV Assay. Anti-HIV screening of test compounds against various viral isolates and cell lines was performed as previously described. (Buckheit et al, Antiviral Res. 1995, 26, 117-132) This cell-based microliter assay quantitates the drug-
 10 induced protection from the cytopathic effect of HIV. Data are presented as the percent control of XTT values for the uninfected, drug-free control. EC_{50} values reflect the drug concentration that provides 50% protection from the cytopathic effect of HIV-1 in infected cultures, while the IC_{50} reflects the concentration of drug that causes 50% cell death in the uninfected cultures. XTT-based results were confirmed by measurement of
 15 cell-free supernatant reverse transcriptase and p24 levels.

Mechanistic Assays. The effects of inhibitors on the *in vitro* activity of purified RT (kind gift of Steve Hughes, NCI-FCRDC, Frederick, MD) were determined by measurement of incorporation of [^{32}P]TTP into the poly(rA):oligo(dT) (rAdT) homopolymer or [^{32}P]GTP into the poly(rC):oligo(dG) (rCdG) template/primer systems.
 20 Samples (5 μL) were blotted onto DE81 paper, washed with 5% diacidic sodium phosphate, and then quantitated on a Packard Matrix 9600 direct beta counter. 3'-Azido-2',3'-dideoxythymidine-5'-triphosphate (AZTTP) and NSC 629243 (UC38) served as a positive control for inhibition of RT.

To determine if compounds affected the HIV-1 nucleocapsid protein zinc
 25 fingers, fluorescence measurements of the Trp 37 residue in the C-terminal zinc finger of the HIV-1 nucleocapsid protein were performed as previously described. (Rice et al. Science, 1995, 270, 1194-1197). The recombinant nucleocapsid protein was prepared at 20 $\mu\text{g/mL}$ in 10 mM sodium phosphate buffer (pH 7.0), treated with 25 μM of test compound, then after indicated time intervals the samples were diluted 1/10 in 10 mM
 30 sodium phosphate buffer (pH 7.0) and the fluorescence intensity measured. The excitation and emission wavelengths utilized with the Shimadzu RF5000 spectrofluorimeter were 280 and 351 nm, respectively. The analytical procedure


employed to determine the reagent-induced inhibition of HIV-1 protease activity has been previously described. Recombinant HIV-1 protease (Bachem BioScience Inc., King of Prussia, PA) and the substrate (Val-Ser-Gln-Asn-Tyr-Pro-Ile-Val-Gln-NH₂, Multiple Peptide Systems, San Diego, CA) were utilized to determine the concentration of test compound required to inhibit protease activity by 50% (IC₅₀). Briefly, HIV-1 protease (14.2 nM final) was mixed with various concentrations of test compounds in 250 mM potassium phosphate buffer, pH 6.5, 2.5% (v/v) glycerol, 0.5 mM dithiothreitol, 0.5 mM EDTA and 375 mM ammonium sulfate, after which the substrate was added (30 nmol) and the reaction incubated at 37 °C for 30 min. Reactions were terminated by the addition of 20 pL of a mixture of 8 M guanidine-HCl to 10% trifluoroacetic acid (8:1), and the reaction products were separated by reverse-phase HPLC on a Nova-Pak C-18 column. Absorbance was measured at 206 nm, peak areas quantitated, and the percentage conversion to product used to calculate the percentage of control cleavage in the presence of inhibitors. The 3'-cleavage and integration activities of purified HIV-1 integrase were quantitated as previously described. (Rice PNAS 1993, 90, 9721-9724).

The attachment/fusion assay was performed as described by Ciminale (AIDS Res. Hum. Retrovir. 1990, 6, 1281-1287) with modification. Briefly, HIV-1 envelope-expressing, Tat-producing HL2/3 cells and CXCR-4 expressing, LTR-βGal-containing MAGI cells (obtained from the AIDS Research and Reference Program, National Institute of Allergy and Infectious Disease, NIH, Bethesda, MD, USA) were preincubated separately with test compound for 1 h at 37°C, followed by admixture of the two cell lines at a cell ratio of one to one. Incubations were then continued for 16 h. The cells were then fixed and stained for the expression of β-gal with indolyl-β-D-galactopyranoside (X-Gal). The number of blue cells (indicating completion of attachment and fusion of membranes) were counted by light microscopy.

Claims:


1. A compound of the formula:

5



wherein X is selected from the group consisting of

10



- 15 wherein R_1 and R_6 are H or halo;

R_2 and R_5 are independently OR_{11} ;


R_3 and R_4 are independently CO_2R_{12} ;

R_8 is selected from the group consisting of $(CH_2)_mOR_{13}$, $(CH_2)_mN_3$, $(CH_2)_mCOOR_{14}$ and $(CH_2)_mNH_3$;

- 20 R_9 and R_{10} are independently selected from the group consisting of H, (C_1-C_5) alkyl, $(CH_2)_nOR_{13}$, $(CH_2)_nN_3$, $(CH_2)_nCOOR_{14}$ and $(CH_2)_nNH_3$;


R_{11} , R_{12} , R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; m is 1-4 and n is 0-4, with the proviso that when X is

25



R_8 is not $(CH_2)_2OH$.

2. The compound of claim 1 wherein X is



5 R_1 and R_6 are independently Br or Cl;


R_2 and R_5 are each OCH_3 ; and

R_3 and R_4 are each CO_2CH_3 .


3. The compound of claim 2 wherein R_8 is $(\text{CH}_2)_m\text{N}_3$ or $(\text{CH}_2)_m\text{COOR}_{14}$.

4. The compound of claim 3 wherein R_{14} is CH_3 or H.

10 5. A compound of the formula:



wherein X is selected from the group consisting of



wherein R_2 and R_5 are independently OR_{11} ;


R_3 and R_4 are independently CO_2R_{12} ;

25 R_7 and R_8 are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;

R_9 and R_{10} are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_n\text{OR}_{13}$, $(\text{CH}_2)_n\text{N}_3$, $(\text{CH}_2)_n\text{COOR}_{14}$ and $(\text{CH}_2)_n\text{NH}_3$;

30 R_{11} , R_{12} , R_{13} and R_{14} are independently selected from the group consisting of H and $(\text{C}_1\text{-C}_5)$ alkyl; m is 1-3 and n is 0-4.

6. The compound of claim 5 wherein X is



5


R_2 and R_3 are each OCH_3 ; and

R_3 and R_4 are each CO_2CH_3 .


7. The compound of claim 6 wherein R_8 is $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$ or $(\text{CH}_2)_m\text{COOR}_{14}$.

10 8. The compound of claim 7 wherein R_{13} is CH_3 or H and R_{14} is CH_3 or H.

9. A composition comprising a reverse transcriptase inhibitory effective amount of a compound of the formula:



wherein X is selected from the group consisting of



wherein R_1 and R_6 are H or halo;

R_2 and R_5 are independently OR_{11} ;

25 R_3 and R_4 are independently CO_2R_{12} ;

R_7 and R_8 are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;

R_9 and R_{10} are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_n\text{OR}_{13}$, $(\text{CH}_2)_n\text{N}_3$, $(\text{CH}_2)_n\text{COOR}_{14}$ and $(\text{CH}_2)_n\text{NH}_3$;

30 R_{11} , R_{12} , R_{13} and R_{14} are independently selected from the group consisting of H and $(\text{C}_1\text{-C}_5)$ alkyl; m is 1-3 and n is 0-4;


and a pharmaceutically acceptable carrier.

10. The composition of claim 9 wherein X is




R_1 and R_6 are independently Br or Cl, R_2 and R_5 are each OCH_3 , R_3 and R_4 are each CO_2CH_3 and R_8 is $(\text{CH}_2)_n\text{OH}$, $(\text{CH}_2)_n\text{COOCH}_3$ or $(\text{CH}_2)_n\text{N}_3$.

10 11. A method of treating a warm-blooded vertebrate suffering from a disease of viral origin, said method comprising the step of administering to said vertebrate a therapeutically effective amount of an antiviral composition comprising a compound of the formula:



wherein X is selected from the group consisting of



wherein R_1 and R_6 are H or halo;

25 R_2 and R_5 are independently OR_{11} ;

R_3 and R_4 are independently CO_2R_{12} ;

R_7 and R_8 are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_m\text{OR}_{13}$, $(\text{CH}_2)_m\text{N}_3$, $(\text{CH}_2)_m\text{COOR}_{14}$ and $(\text{CH}_2)_m\text{NH}_3$;

30 R_9 and R_{10} are independently selected from the group consisting of H, $(\text{C}_1\text{-C}_5)$ alkyl, $(\text{CH}_2)_n\text{OR}_{13}$, $(\text{CH}_2)_n\text{N}_3$, $(\text{CH}_2)_n\text{COOR}_{14}$ and $(\text{CH}_2)_n\text{NH}_3$;


R_{11} and R_{12} are independently $(\text{C}_1\text{-C}_5)$ alkyl or H;

R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl;
m is 1-3 and n is 0-4;

and a pharmaceutically acceptable carrier.

12. The composition of claim 11 wherein X is

5



10 R_1 and R_2 are independently Br or Cl;

R_2 and R_5 are each OCH_3 ; and

R_3 and R_4 are each CO_2CH_3 .

13. The composition of claim 12 wherein R_8 is $(CH_2)_mOR_{13}$, $(CH_2)_mN_3$ or $(CH_2)_mCOOR_{14}$.

15

Abstract: A novel alkenyldiarylmethane (ADAM) analog has been synthesized with enhanced potency as an anti-HIV agent. The new compound (ADAM II) inhibits the cytopathic effect of HIV-1_{RF} in CEM-SS cells with an EC₅₀ of 31.6 μ M, providing a therapeutic index of 158. ADAM II is a non-nucleoside reverse transcriptase inhibitor,
5 displaying an IC₅₀ of 0.5 μ M with poly(rC)oligo(dT) as the template/primer.

PROVISIONAL
PATENT APPLICATION
of
Mark Cushman
Agustin Casimiro-Garcia
William G Rice
for
**ALKENYLDIARYLMETHANE NON-NUCLEOSIDE HIV-1
REVERSE TRANSCRIPTASE INHIBITORS**

Attorney Docket 3220-29595

Attorneys:

John P. Breen
Steven R. Lammert
BARNES & THORNBURG
11 South Meridian Street
Indianapolis, Indiana 46204

Mark Cushman, *[†] Agustin Casimiro-Garcia,[†] Karen Williamson,[†] and William G. Rice[†]

It was decided to synthesize an ADAM analog incorporating a methyl ester at the end of the chain.

The desired analog ADAM II (3) was prepared in one step from known starting materials. Reaction of substituted benzophenone 4⁷ with the Wittig reagent 5¹⁰ in THF at -78 °C for 12 hours followed by room temperature for 12 hours afforded the desired compound 3 in 45% isolated yield as a pale yellow oil after silica gel chromatography, eluting with hexane-ethyl acetate 3:1.

ADAM II (3) was tested for prevention of the cytopathic effect of HIV-1_{RF} in CEM-SS cells. The EC₅₀ for prevention of the cytopathic effect of the virus in cell culture was 0.2 μM, while the IC₅₀ for cytotoxicity in uninfected cells was 31.6 μM. This corresponds to a therapeutic index (IC₅₀/EC₅₀) of 158. ADAM II was also tested as an inhibitor of HIV-1 reverse transcriptase. With poly(rC)oligo(dT) as the template/primer, ADAM II inhibited the enzyme with an IC₅₀ of 0.5 μM, while with poly(rA)oligo(dT) as the template/primer, the compound exerted an IC₅₀ of 1.9 μM. The greater sensitivity to inhibition with poly(rA)oligo(dT) as the template primer is characteristic of the non-nucleoside HIV-1 reverse transcriptase inhibitors.¹¹⁻¹⁶ In order to confirm that ADAM II (3) was indeed acting as an NNRTI, it was tested in a number of assays representative of important events in the replication of HIV-1. The results indicated that ADAM II (3) did not have any significant effect on integrase, protease, or nucleocapsid protein zinc fingers.

There is a significant difference in potency between 1 and 3 for prevention of the cytopathic effect of HIV-1 in cell culture. For ADAM 1, the EC₅₀ for the prevention of the cytopathic effect of HIV-1_{RF} in CEM-SS cells proved to be 9.2 μM, while the IC₅₀ for cytotoxicity in uninfected CEM-SS cells was 138 μM. This corresponds to a therapeutic index (IC₅₀/EC₅₀) of only 15. The anti-HIV potency of ADAM II (3) is greater than that of 1 by a factor of 46. The terminal methoxycarbonyl group present on the side chain of ADAM

II (3) may in fact be acting as a hydrogen bond acceptor from Glu 138 or Lys 103. However, a definitive answer to this question would have to await the crystal structure of the enzyme complexed with the bound ligand. The difference in activity is most likely not due to the dichloro substitution in 3 vs. the dibromo substitution in 1, since the analog of ADAM in which the two bromines are replaced by chlorines is less active than 1 vs. HIV-1_{RF} in CEM cells.

The orientations of the two aromatic systems in 1 and 3 fit a "butterfly-like" model recently proposed on the basis of a comparison of the structures of several non-nucleoside HIV-1 reverse transcriptase inhibitors.¹⁷ The hexenyl side chain of 1 points toward a "lipophilic site" proposed by the model.

Future work will concentrate on defining the resistance mutations to ADAM II and also investigating the anti-HIV activity of ADAM II in animal models.

Acknowledgment. This research was made possible by NIH Grant NO1-AI-36624. A. C.-G. wishes to thank Consejo Nacional de Ciencia y Tecnologia (Mexico) for a scholarship.

The non-nucleoside HIV-1 reverse transcriptase inhibitors (NNRTIs) comprise a structurally diverse set of compounds that inhibit the enzyme by an allosteric mechanism involving binding to a site adjacent to the deoxyribonucleoside triphosphate binding site.¹ Well known NNRTIs include hydroxethoxymethylphenylthiothymine (HEPT),² tetrahydroimidazobenzodiazepinone (TIBO),³ dipyrindodiazepinone (nevirapine),⁴ pyridinone,⁵ bis(heteroaryl)piperazine (BHAP),⁶ *tert*-butyldimethylsilylspiroaminooxathiole dioxide (TSAO),⁷ and α -anilinophenylacetamide (α -APA) derivatives. Although the rapid emergence of resistant viral strains has hampered the clinical development of the NNRTIs for the treatment of AIDS, several strategies have emerged for overcoming resistance, including switching to another NNRTI to which the virus has remained sensitive, using higher doses of the NNRTI against the resistant strain,^{9,10} and employment of combinations of agents which elicit mutations that counteract one another.^{11,12} A need remains for additional NNRTIs having unique patterns of resistance mutations in order to facilitate the application of these strategies.


We recently reported a series of alkenyldiarylmethanes (ADAMs) which act as NNRTIs.¹³ The most potent of these proved to be ADAM 1, which displayed synergistic anti-HIV-1 activity with AZT and was found to be more effective against an AZT-resistant

virus. However, the potency range of ADAM 1 (0.56 to 16 μ M) against a variety of non-resistant HIV-1 strains was lower than that generally observed with many of the known NNRTIs. In order to improve on the lead provided by ADAM 1, additional compounds in this series were prepared and evaluated both as inhibitors of HIV-1 reverse transcriptase (RT) and for inhibition of the cytopathic effect of HIV in cell culture. As described below, this has resulted in the design and synthesis of a very potent alkenyldiarylmethane which inhibits the cytopathic effect of HIV-1_{RF} in CEM-SS cells with an EC_{50} of 13 nM, while it shows cytotoxicity with an IC_{50} of 31.6 μ M, providing a therapeutic index of 2430.


To facilitate the design of novel alkenyldiarylmethane compounds that exhibit inhibitory reverse transcriptase activity, a hypothetical model was constructed of the binding of ADAM I (1) to HIV-I reverse transcriptase. This model was derived by overlaying the structure of ADAM I (1) "on top" of the ligand in the crystal structures of nevirapine (2) bound to HIV-1 reverse transcriptase. The structure of nevirapine was then removed, the protein was "frozen", and the energy of the resulting ADAM-RT complex was minimized while the ligand was allowed to move. The validity of this procedure is supported by the fact nevirapine, α -APA, and TIBO all bind to HIV-1 in a nearly identical mode which conforms to a general "butterfly" structural model for NNRTIs.^{8,9} **NEED REFERENCE CITE** The model is also supported by mutagenesis studies of the ADAM binding site on HIV-1 reverse transcriptase, in which it was determined that the resistance mutations to ADAM I (1) circumscribe a well-defined binding pocket.⁶ According to the model, the end of the side chain of ADAM I (1) is located near Glu 138 and Lys 103 of HIV-1 reverse transcriptase. Both of these residues contain bond donors. It might be expected that the incorporation of functional groups at the end of the alkenyl chain of the ligand which are capable of acting as hydrogen bond acceptors might interact favorably with the terminal amino group of Lys 103 or the terminal carboxylic acid group of Glu 138.

The present invention is directed to non-nucleoside compounds that inhibit reverse transcriptase activity, their pharmaceutical compositions and methods utilizing such compounds/compositions for treating patients suffering from a viral infection. More particularly the compounds of the present invention are useful for treating patients suffering from a disease of retroviral origin, such as HIV.

The compounds of the present invention are alkenyldiarylmethane compounds of formula I:



wherein X is selected from the group consisting of



wherein R_1 and R_6 are H or halo;

R_2 and R_5 are independently OR_{11} ;


R_3 and R_4 are independently CO_2R_{12} ;

R_7 , R_8 , R_9 and R_{10} are independently selected from the group consisting of H, (C_1-C_5) alkyl, $(CH_2)_nOR_{13}$, $(CH_2)_nN_3$, $(CH_2)_nCOOR_{14}$, $(CH_2)_nOSO_2CH_3$ and $(CH_2)_nNH_3$;

R_{11} and R_{12} are independently (C_1-C_3) alkyl or H;


R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; and n is 0-4.

In one preferred embodiment of this invention there is provided compounds of the above formula I, wherein X is



R_1 and R_6 are independently Br or Cl, R_2 and R_5 are each OCH_3 , R_3 and R_4 are each CO_2CH_3 , R_7 is H and R_8 is $(CH_2)_mOH$, $(CH_2)_mCOOCH_3$ or $(CH_2)_mN_3$, wherein m is 2 or 3. These compounds have been demonstrated to inhibit the cytopathic effect of HIV-1_{RF} in CEM-SS cells. (See Table 1).

In another preferred embodiment of this invention there is provided a reverse transcriptase inhibiting compounds of the above formula I wherein X is



R_1 and R_6 are halo;

R_2 and R_5 are OCH_3 ;

R_3 and R_4 are CO_2CH_3 ;


R_7 is H;

R_8 is (C_2-C_5) alkyl, $(CH_2)_mOR_{13}$, $(CH_2)_mN_3$, $(CH_2)_mCOOR_{14}$ and $(CH_2)_mNH_3$;

R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; and m is 2-3.


The compounds of this invention are readily formulated into pharmaceutical compositions, also within the scope of this invention, for use in the presently described method for treatment of patients suffering from a disease of retroviral origin. In one preferred embodiment of this invention, the pharmaceutical composition comprises a reverse transcriptase inhibitory effective amount of an compound of formula I:

I




wherein X is selected from the group

consisting of



and



wherein R_1 and R_6 are H or halo;

R_2 and R_3 are independently OR_{11} ;

R_3 and R_4 are independently CO_2R_{12} ;


R_7 and R_8 are independently selected from the group consisting of H, (C_1-C_5) alkyl, $(CH_2)_mOR_{13}$, $(CH_2)_mN_3$, $(CH_2)_mCOOR_{14}$ and $(CH_2)_mNH_3$;

R_9 and R_{10} are independently selected from the group consisting of H, (C_1-C_5) alkyl, $(CH_2)_nOR_{13}$, $(CH_2)_nN_3$, $(CH_2)_nCOOR_{14}$ and $(CH_2)_nNH_3$;

R_{11} and R_{12} are independently (C_1-C_5) alkyl or H;

R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; m is 1-3 and n is 0-4 and a pharmaceutically acceptable carrier.

Another pharmaceutical composition within the scope of this invention comprises a reverse transcriptase inhibiting compounds of the above formula I wherein X is



R_1 and R_6 are halo;

R_2 and R_5 are OCH_3 ;

R_3 and R_4 are CO_2CH_3 ;

R_8 is (C_2-C_5) alkyl, $(CH_2)_mOR_{13}$, $(CH_2)_mN_3$, $(CH_2)_mCOOR_{14}$ and $(CH_2)_mNH_3$;

R_{13} and R_{14} are independently selected from the group consisting of H and (C_1-C_5) alkyl; and m is 2-3 and a pharmaceutically acceptable carrier.

The anti-viral activity of the described compounds was measured utilizing two different assays. The first assay measures the effectiveness of the alkenyldiarylmethane compound to inhibit reverse transcriptase activity, and the second assay measured the compound's ability to inhibit the cytopathic activity of HIV-1 to cells cultured *in vitro*.

The present invention further provides pharmaceutical formulations comprising an effective amount of an alkenyldiarylmethane compound for treating a patient suffering from a disease of retroviral origin. As used herein, an effective amount of the alkenyldiarylmethane compound is defined as the amount of the compound which, upon

administration to a patient, alleviates or eliminates symptoms of the disease, or reduces or eliminates detectable levels of the virus in the treated patient.

The effective amount to be administered to a patient is typically based on body surface area, patient weight, and patient condition. Body surface area may be approximately determined from patient height and weight (see e.g., Scientific Tables, Geigy Pharmaceuticals, Ardley, New York, pages 537-538 (1970)). Effective doses will vary, as recognized by those skilled in the art, dependant on route of administration, excipient usage and the possibility of co-usage with other therapeutic treatments including other anti-tumor agents, and radiation therapy.

The pharmaceutical formulation may be administered via the parenteral route, including subcutaneously, intraperitoneally, intramuscularly and intravenously. Examples of parenteral dosage forms include aqueous solutions of the active agent, in a isotonic saline, 5% glucose or other well-known pharmaceutically acceptable liquid carrier. In one preferred aspect of the present embodiment, the alkenyldiarylmethane compound is dissolved in a saline solution containing 5% of dimethyl sulfoxide and 10% Cremphor EL (Sigma Chemical Company). Additional solubilizing agents well-known to those familiar with the art, can be utilized as pharmaceutical excipients for delivery of the selenophene compounds.

The present compound can also be formulated into dosage forms for other routes of administration utilizing well-known methods. The pharmaceutical compositions can be formulated, for example, in dosage forms for oral administration in a capsule, a gel seal or a tablet. Capsules may comprise any well-known pharmaceutically acceptable material such as gelatin or cellulose derivatives. Tablets may be formulated in accordance with conventional procedure by compressing mixtures of the active polythiophene and solid carriers, and lubricants well-known to those familiar with the art. Examples of solid carriers include starch, sugar, bentonite. The compounds of the present invention can also be administered in a form of a hard shell tablet or capsule containing, for example, lactose or mannitol as a binder and conventional fillers and tableting agents.

The following examples are provided to illustrate various embodiments of Applicants' invention, and are not intended to in any way limit the scope of the invention as set forth in this specification and appended claims.

Compound	RT (I_{50} μ M)		XTT Assay		
	rCdG	rAdT	EC ₅₀	IC ₅₀	TI ₅₀
Bfly	10.1	> 100	NI ₁₀₀	> 100	-
Bfly-Et	14	> 100	NI ₁₀₀	> 100	-
Bfly-OMe	96	> 100	-	26	-
EH-4-29	5.6	> 100	NI ₁₀₀	> 100	-
EH-5-21	> 100				
EH-5-23	26		-	10	-
EH-5-57	100		-	40	-
EH-5-66	31.6		6.0	> 316	> 52
EH-5-68	94		1.1	> 316	> 287
EH-5-70	> 100		-	22	-
FM-3-1					
FM-3-4	> 100	> 100			
JR-F-27B	3.2		-	14	-
JR-F-57	98		-	57	-
JR-E-76B	11		-	> 316	-
ACG-8602-72	> 100	> 100			
ACG-8752-17A	> 100	> 100			
ACG-8752-43	> 100	> 100			
ACG-8752-53	> 100	> 100			
ACG-8752-64					
ACG-8752-80			-	6.9	-
ACG-8752-15			-	176	-
ACG-8825-4			0.6	35	58
ACG-8825-9			NI ₁₀₀	> 100	-
ACG-8825-31	0.5		0.013	31.6	2430
ACG-8825-34	63		-	14.9	-
ACG-8825-48	> 100	> 100	-	48.0	-
ACG-8825-50	> 100	> 100	-	38.8	-
ACG-8825-54	> 100	> 100	-	61.2	-
ACG-8825-59			-	24.9	-

Supplier: Mark Cushman (055W)

Date: 4 Apr 96


Sample No: Bfly

Amount Supplied: 20 mg

Name: 6H,12H-6,12-Methyliminedibenzo[b,f][1,5]dioxocin

Structural Formula:

NSC # D687941-U



Molecular Formula: $C_{15}H_{13}NO_2$

Parent Analog:

MW: 238

Target Compound: .

Stability:

Solubility: $CHCl_3$

Other Data: NMR.

45

Supplier: Mark Cushman (055W)

Date: 4 Apr 96


Sample No: Bfly-Et

Amount Supplied: 20 mg

Name: 6H,12H-6,12-Ethyliminedibenzo[b,f][1,5]dioxocin

Structural Formula:

NSC # D687940-T



Molecular Formula: C₁₆H₁₅NO₂

Parent Analog:

MW: 253.3

Target Compound: .

Stability:

Solubility: CHCl₃

Other Data: IR, MS, elemental, mp=131-2°C, ¹H NMR.

Supplier: Mark Cushman (055W)

Date: 4 May 96


Sample No: Bfly-OMe

Amount Supplied: 20 mg

Name: 6H,12H-4,10-Dimethoxy-6,12-methyliminedibenzo[b,f][1,5]dioxocin

Structural Formula:

NSC # D687942-V



Molecular Formula: C₁₇H₁₇NO₄

Parent Analog:

MW: 299.33


Target Compound:

Stability:

Solubility: CHCl₃

Other Data: MS, Elemental, mp=208-210°C, ¹H NMR.


NSC 694754

DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 01/14/97
Supplier Compound ID: EH-5-70
NSC Number: 694754-Z
Molecular Formula: C₂₁H₂₀Br₂O₇
Acquired for Testing Against: AIDS

NSC 694556




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 01/10/97
Supplier Compound ID: EE-5-68
NSC Number: 694556-L
Molecular Formula: C22H21Br2N3O6
Acquired for Testing Against: AIDS

NSC 694249




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 12/13/96
Supplier Compound ID: EH-5-66
NSC Number: 694249-A
Molecular Formula: C₂₂H₂₂Br₂O₇
Acquired for Testing Against: AIDS

NSC 694248




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 12/13/96
Supplier Compound ID: EH-5-57
NSC Number: 694248-Z
Molecular Formula: C₂₁H₂₂
Acquired for Testing Against: AIDS

NSC 694555




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 01/10/97
Supplier Compound ID: EH-5-23
NSC Number: 694555-K
Molecular Formula: C₂₁H₂₀Br₂O₆
Acquired for Testing Against: AIDS

NSC 694251




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 12/13/96
Supplier Compound ID: EH-5-21
NSC Number: 694251-F
Molecular Formula: C₂₀H₁₈Br₂O₆
Acquired for Testing Against: AIDS

NSC 686377



DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCT, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 03/18/96
Supplier Compound ID: EH - 4-29
NSC Number: 686377-U
Molecular Formula: C₂₅ H₂₈ Br₂ O₇
Acquired for Testing Against: AIDS

Supplier: Mark Cushman (055W)

Sample No: FM-3-1


Amount Supplied 20 mg

Date 09.13.96

Name: 5-[4-(Fluorosulfonyl)benzoylamino]-6-ribitylamino uracil

Structural Formula

NSC # 691615-0



Molecular Formula $C_{16}H_{19}N_4O_9SF$

Parent Analog

MW 462 (corrected* : 506.53)

Target Compound

Stability Unstable in aqueous base


Solubility H_2O

Other Data Anal. : Calcd for $C_{16}H_{19}N_4O_9SF \cdot 0.2NaCl \cdot 1.8H_2O$: C, 37.94; H, 4.50; N, 11.06; S, 6.33; F, 3.75. Found: C, 37.58; H, 4.18; N, 11.11; S, 6.16; F, 3.56.

MS, NMR.

* includes solvents found upon elemental analysis.

55
NSC 692220




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 10/04/96
Supplier Compound ID: FM-3-4
NSC Number: 692220-V
Molecular Formula: C₁₁H₈FN₃O₅S
Acquired for Testing Against: AIDS

NSC 694065




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 12/09/96
Supplier Compound ID: JR-F-27
NSC Number: 694065-A
Molecular Formula: C₂₅H₃₀O₆
Acquired for Testing Against: AIDS

NSC 694064



DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 12/09/96
Supplier Compound ID: JR-E-76
NSC Number: 694064-Z
Molecular Formula: C₂₅H₂₈I₂O₆
Acquired for Testing Against: AIDS

Supplier: Mark Cushman

Date July 18th, 1996


Sample No: ACG-8602-72

Amount Supplied: 20 mg

Name: 4,4-Bis[8',8''-dichloro-2',2',2'',2''-tetramethyl-4',4''-dioxo-6',6''-(1,3-benzodioxyl)]-3-buten-1-ol

Structural Formula:

690046-J .



Molecular Formula: $C_{24}H_{22}Cl_2O_7$


MW: 493.3447

Stability: Stable

Solubility: Chloroform, DMSO, acetone

Other Data: Elemental analysis, IR, MS, NMR, m.p. 185-188 °C

NSC 690985




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 08/22/96
Supplier Compound ID: ACG-8752-43
NSC Number: 690985-F
Molecular Formula: C₂₂H₂₂Cl₂O₇
Acquired for Testing Against: AIDS

NSC 690986

DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 08/22/96
Supplier Compound ID: ACG-8752-53
NSC Number: 690986-G
Molecular Formula: C₂₁H₂₀Cl₂O₆
Acquired for Testing Against: AIDS

Supplier: Mark Cushman

Date September 13, 1996


Sample No: ACG-8752-64

Amount Supplied: 20 mg

Name: 4,4-(3',3''-dicarbomethoxy-5',5''-dichloro-4',4''-dimethoxydiphenyl)-3-butenic acid

Structural Formula:

691616-P



Molecular Formula: C₂₂H₂₀Cl₂O₆


MW: 483.3026

Stability: Stable solid

Solubility: Acetone, ethyl ether, DMSO.

Other Data: IR, NMR, m.p.: 206-207 °C

NSC 692623



DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 10/15/96
Supplier Compound ID: ACG-8752-80
NSC Number: 692623-K
Molecular Formula: C₂₃H₂₄Cl₂O₉S
Acquired for Testing Against: AIDS

Supplier: Mark Cushman


Date October 24, 1996

Sample No: ACG-8752-15

Amount Supplied: 20 mg

Name: 4-Bromo-1,1-bis[8',8''-dichloro-2',2',2'',2''-tetramethyl-4',4''-dioxo-6',6''-(1,3-benzodioxyl)]-1-butene

Structural Formula:



235-7

692751-X

offshot P

Molecular Formula: $C_{24}H_{21}BrCl_2O_6$


MW: 553.9898

Stability: Pale yellow solid

Solubility: Chloroform, acetone, DMSO.

Other Data: IR, NMR, MS, Elemental Analysis, m.p.: 176-178 °C

64
NSC 692624



DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 10/15/96
Supplier Compound ID: ACG-8825-4
NSC Number: 692624-L
Molecular Formula: C₂₂H₂₁Cl₂N₃O₆
Acquired for Testing Against: AIDS

Supplier: Mark Cushman


Date October 24, 1996

Sample No: ACG-8825-9

Amount Supplied: 20 mg

Name: *tert*-Butyl 3,3-(3',3''-Dicarbomethoxy-5',5''-dichloro-4',4''-dimethoxydiphenyl)-2-propenoate

Structural Formula:



236-U

692752-Y

whf

Molecular Formula: C₂₅H₂₆Cl₂O₅


MW: 524.1004

Stability: White solid

Solubility: Chloroform, acetone, DMSO.

Other Data: IR, NMR, MS, m.p.: 114-115.5 °C

NSC 695574




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 02/13/97
Supplier Compound ID: ACG-8825-31
NSC Number: 695574-R
Molecular Formula: C₂₅H₂₆Cl₂O₈
Acquired for Testing Against: AIDS

67

NSC 695573




DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 02/13/97
Supplier Compound ID: ACG-8825-34
NSC Number: 695573-Q
Molecular Formula: C₂₂H₂₃Cl₂NO₆.HCl
Acquired for Testing Against: AIDS


NSC 696043

DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 03/05/97
Supplier Compound ID: ACG-8825-48
NSC Number: 696043-A
Molecular Formula: C₂₁H₁₈Cl₂O₈
Acquired for Testing Against: AIDS


NSC 696221

DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 03/17/97
Supplier Compound ID: ACG-8825-54
NSC Number: 696221-U
Molecular Formula: C₄₈H₆₅Cl₂N₁O₇
Acquired for Testing Against: AIDS

NSC 696443



DRUG SYNTHESIS AND CHEMISTRY BRANCH, DTP, DCTDC, NCI
NSC NUMBER LIST

Dr. Mark Cushman
Dept of Medicinal Chemistry & Pharmacognosy
School of Pharmacy and Pharmacal Sciences
Purdue University
1333 Robert E. Heine Pharmacy Building
West Lafayette, IN 47907-1333

Transmittal Letter or Collection Date: 03/26/97
Supplier Compound ID: ACG-8825-59
NSC Number: 696443-M
Molecular Formula: C₂₂H₂₀Cl₂O₈
Acquired for Testing Against: AIDS